

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>C12Q 1/68, C12P 19/34, C07H 21/04</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 99/09217</b> <b>(43) International Publication Date:</b> 25 February 1999 (25.02.99)
<b>(21) International Application Number:</b> PCT/US98/16966 <b>(22) International Filing Date:</b> 14 August 1998 (14.08.98)  <b>(30) Priority Data:</b> 08/912,885      15 August 1997 (15.08.97)      US 08/947,779      9 October 1997 (09.10.97)      US 08/959,365      28 October 1997 (28.10.97)      US  <b>(71) Applicant (for all designated States except US):</b> HYSEQ, INC. [US/US]; 670 Almanor Avenue, Sunnyvale, CA 94086 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> DRMANAC, Radoje [YU/US]; 850 East Greenwich Place, Palo Alto, CA 94303 (US). DRMANAC, Snezana [YU/US]; 850 East Greenwich Place, Palo Alto, CA 94303 (US). BAIDYA, Narayan [IN/US]; 966 Helen Avenue #1, Sunnyvale, CA 94086 (US).  <b>(74) Agents:</b> ABRAMS, Samuel, B. et al.; Pennie & Edmonds LLP, 1155 Avenue of the Americas, New York, NY 10036 (US).		<b>(81) Designated States:</b> AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, HR, HU, ID, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> METHODS AND COMPOSITIONS FOR DETECTION OR QUANTIFICATION OF NUCLEIC ACID SPECIES		
<b>(57) Abstract</b> <p>The present invention provides a method for detecting a target nucleic acid species using an array of probes affixed to a substrate and a plurality of labeled probes. The invention also relates to oligonucleotide probes attached to discrete particles wherein the particles can be grouped into a plurality of sets based on a physical property. A different probe is attached to the discrete particles of each set, and the identity of the probe is determined by identifying the discrete particles from their physical property. The invention further relates to methods using agents which destabilize the binding of complementary polynucleotide strands (decrease the binding energy) or increase stability of binding between complementary polynucleotide strands (increase the binding energy). The figure is an illustration of an apparatus for mass producing probe arrays.</p> <div data-bbox="844 1155 1315 1722"> </div>		

BEST AVAILABLE COPY

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

WO 99/09217

PCT/US98/16966

## METHODS AND COMPOSITIONS FOR DETECTION OR QUANTIFICATION OF NUCLEIC ACID SPECIES

### CROSS REFERENCE TO RELATED APPLICATIONS

- 5 This patent application is a continuation-in-part of U.S. Patent application Serial No. 08/912,885, filed on August 15, 1997, and a continuation-in-part of U.S. Patent application Serial No. 08/947,779, filed on October 9, 1997, and a continuation-in-part of U.S. Patent application Serial No. 08/892,503, filed on July 14, 1997, and a continuation-in-part of U.S. Patent application Serial No. 08/812,951, filed on March 4, 10 1997, and a continuation-in-part of U.S. Patent application Serial No. 08/784,747, filed on January 16, 1997.

### FIELD OF THE INVENTION

- This invention relates in general to methods and apparatus for nucleic acid 15 analysis, and, in particular, to methods and apparatus for nucleic acid analysis.

### BACKGROUND

- The rate of determining the sequence of the four nucleotides in nucleic acid samples is a major technical obstacle for further advancement of molecular biology, 20 medicine, and biotechnology. Nucleic acid sequencing methods which involve separation of nucleic acid molecules in a gel have been in use since 1978. The other proven method for sequencing nucleic acids is sequencing by hybridization (SBH).

- The traditional method of determining a sequence of nucleotides ( i.e., the order of the A, G, C and T nucleotides in a sample) is performed by preparing a mixture of 25 randomly-terminated, differentially labelled nucleic acid fragments by degradation at specific nucleotides, or by dideoxy chain termination of replicating strands. Resulting nucleic acid fragments in the range of 1 to 500 bp are then separated on a gel to produce a ladder of bands wherein the adjacent samples differ in length by one nucleotide.

- The array-based approach of SBH does not require single base resolution in 30 separation, degradation, synthesis or imaging of a nucleic acid molecule. Using mismatch discriminative hybridization of short oligonucleotides K bases in length, lists of constituent K-mer oligonucleotides may be determined for target nucleic acid. Sequence

WO 99/09217

PCT/US98/16966

for the target nucleic acid may be assembled by uniquely overlapping scored oligonucleotides.

There are several approaches available to achieve sequencing by hybridization. In a process called SBH Format 1, nucleic acid samples are arrayed, and labeled probes are hybridized with the samples. Replica membranes with the same sets of sample nucleic acids may be used for parallel scoring of several probes and/or probes may be multiplexed. Nucleic acid samples may be arrayed and hybridized on nylon membranes or other suitable supports. Each membrane array may be reused many times. Format 1 is especially efficient for batch processing large numbers of samples.

10 In SBH Format 2, probes are arrayed at locations on a substrate which correspond to their respective sequences, and a labelled nucleic acid sample fragment is hybridized to the arrayed probes. In this case, sequence information about a fragment may be determined in a simultaneous hybridization reaction with all of the arrayed probes. For sequencing other nucleic acid fragments, the same oligonucleotide array may be reused.

15 The arrays may be produced by spotting or by in situ synthesis of probes.

In Format 3 SBH, two sets of probes are used. In one embodiment, a set may be in the form of arrays of probes with known positions, and another, labelled set may be stored in multiwell plates. In this case, target nucleic acid need not be labelled. Target nucleic acid and one or more labelled probes are added to the arrayed sets of probes. If

20 one attached probe and one labelled probe both hybridize contiguously on the target nucleic acid, they are covalently ligated, producing a detected sequence equal to the sum of the length of the ligated probes. The process allows for sequencing long nucleic acid fragments, e.g. a complete bacterial genome, without nucleic acid subcloning in smaller pieces.

25 In the present invention, SBH is applied to the efficient identification and sequencing of one or more nucleic acid samples. The procedure has many applications in nucleic acid diagnostics, forensics, and gene mapping. It also may be used to identify mutations responsible for genetic disorders and other traits, to assess biodiversity and to produce many other types of data dependent on nucleic acid sequence.

30



WO 99/09217

PCT/US98/16966

**SUMMARY OF THE INVENTION**

The present invention provides a method for detecting a target nucleic acid species including the steps of providing an array of probes affixed to a substrate and a plurality of labeled probes wherein each labeled probe is selected to have a first nucleic acid sequence  
5 which is complementary to a first portion of a target nucleic acid and wherein the nucleic acid sequence of at least one probe affixed to the substrate is complementary to a second portion of the nucleic acid sequence of the target, the second portion being adjacent to the first portion; applying a target nucleic acid to the array under suitable conditions for hybridization of probe sequences to complementary sequences; introducing a labeled  
10 probe to the array; hybridizing a probe affixed to the substrate to the target nucleic acid; hybridizing the labeled probe to the target nucleic acid; affixing the labeled probe to an adjacently hybridized probe in the array; and detecting the labeled probe affixed to the probe in the array. According to preferred methods of the invention the array of probes affixed to the substrate comprises a universal set of probes. According to other preferred  
15 aspects of the invention at least two of the probes affixed to the substrate define overlapping sequences of the target nucleic acid sequence and more preferably at least two of the labelled probes define overlapping sequences of the target nucleic acid sequences. Still further, according to another aspect of the invention a method is provided for detecting a target nucleic acid of known sequence comprising the steps of:  
20 contacting a nucleic acid sample with a set of immobilized oligonucleotide probes attached to a solid substrate under hybridizing conditions wherein the immobilized probes are capable of specific hybridization with different portions of said target nucleic acid sequence; contacting the target nucleic acid with a set of labelled oligonucleotide probes in solution under hybridizing conditions wherein the labeled probes are capable of specific  
25 hybridization with different portions of said target nucleic acid sequence adjacent to the immobilized probes; covalently joining the immobilized probes to labelled probes that are immediately adjacent to the immobilized probe on the target sequence (e.g., with ligase); removing any non-ligated labelled probes; detecting the presence of the target nucleic acid by detecting the presence of said labelled probe attached to the immobilized probes. The  
30 invention also provides a method of determining expression of a member of a set of partially or completely sequenced genes in a cell type, a tissue or a tissue mixture comprising the steps of: defining pairs of fixed and labeled probes specific for the

WO 99/09217

PCT/US98/16966

sequenced gene; hybridizing unlabeled nucleic acid sample and corresponding labeled probes to one or more arrays of fixed probes; forming covalent bonds between adjacent hybridized labeled and fixed probes; removing unligated probes; and determining the presence of the sequenced gene by detection of labeled probes bound to prespecified  
5 locations in the array. In a preferred embodiment of this aspect of the invention, the target nucleic acid will identify the presence of an infectious agent.

Further, the present invention provides for an array of oligonucleotide probes comprising a nylon membrane; a plurality of subarrays of oligonucleotide probes on the nylon membrane, the subarrays comprising a plurality of individual spots wherein each  
10 spot is comprised of a plurality of oligonucleotide probes of the same sequence; and a plurality of hydrophobic barriers located between the subarrays on the nylon membrane, whereby the plurality of hydrophobic barriers prevents cross contamination between adjacent subarrays.

Still further, the present invention provides a method for sequencing a repetitive  
15 sequence, having a first end and a second end, in a target nucleic acid comprising the steps of: (a) providing a plurality of spacer oligonucleotides of varying lengths wherein the spacer oligonucleotides comprise the repetitive sequence; (b) providing a first oligonucleotide that is known to be adjacent to the first end of the repetitive sequence; (c) providing a plurality of second oligonucleotides one of which is adjacent to the second  
20 end of the repetitive sequence, wherein the plurality of second oligonucleotides is labeled; (d) hybridizing the first and the plurality of second oligonucleotides, and one of the plurality of spacer oligonucleotides to the target nucleic acid ; (e) ligating the hybridized oligonucleotides; (f) separating ligated oligonucleotides from unligated oligonucleotides; and (g) detecting label in the ligated oligonucleotides.

25 Still further, the present invention provides a method for sequencing a branch point sequence, having a first end and a second end, in a target nucleic acid comprising the steps of: (a) providing a first oligonucleotide that is complementary to a first portion of the branch point sequence wherein the first oligonucleotide extends from the first end of the branch point sequence by at least one nucleotide; (b) providing a plurality of  
30 second oligonucleotides that are labeled, and are complementary to a second portion of the branch point sequence wherein the plurality of second oligonucleotides extend from the second end of the branch point sequence by at least one nucleotide, and wherein the

WO 99/09217

PCT/US98/16966

portion of the second oligonucleotides that extend from the second end of the branch point sequence comprise sequences that are complementary to a plurality of sequences that arise from the branch point sequence; (c) hybridizing the first oligonucleotide, and one of the plurality of second oligonucleotides to the target DNA; (d) ligating the hybridized  
5 oligonucleotides; (e) separating ligated oligonucleotides from unligated oligonucleotides; and (f) detecting label in the ligated oligonucleotides.

Still further, the present invention provides a method for confirming a sequence by using probes that are predicted to be negative for the target nucleic acid. The sequence of a target is then confirmed by hybridizing the target nucleic acid to the "negative" probes  
10 to confirm that these probes do not form perfect matches with the target nucleic acid.

Still further, the present invention provides a method for analyzing a nucleic acid using oligonucleotide probes that are complexed with different labels so that the probes may be multiplexed in a hybridization reaction without a loss of sequence information (i.e., different probes have different labels so that hybridization of the different probes to  
15 the target can be distinguished). In a preferred embodiment, the labels are radioisotopes, or fluorescent molecules, or enzymes, or electrophore mass labels. In a more preferred embodiment, the differently labeled oligonucleotides probes are used in format III SBH, and multiple probes (more than two, with one probe being the immobilized probe) are ligated together.

20 Still further, the present invention provides a method for detecting the presence of a target nucleic acid having a known sequence when the target is present in very small amounts compared to homologous nucleic acids in a sample. In a preferred embodiment, the target nucleic acid is an allele present at very low frequency in a sample that has nucleic acids from a large number of sources. In an alternative preferred embodiment,  
25 the target nucleic acid has a mutated sequence, and is present at very low frequency within a sample of nucleic acids.

Still further, the present invention provides a method for confirming the sequence of a target nucleic acid by using single pass gel sequencing. Primers for single pass gel sequencing are derived from the sequence obtained by SBH, and these primers are used in  
30 standard Sanger sequencing reactions to provide gel sequence information for the target nucleic acid. The sequence obtained by single pass gel sequencing is then compared to the SBH derived sequence to confirm the sequence.

WO 99/09217

PCT/US98/16966

Still further, the present invention provides a method for solving branch points by using single pass gel sequencing. Primers for the single pass gel sequencing reactions are identified from the ends of the Sfs obtained after a first round of SBH sequencing, and these primers are used in standard Sanger-sequencing reactions to provide gel sequencing  
5 information through the branch points of the Sfs. Sfs are then aligned by comparing the Sanger-sequencing results through the branch points to the Sfs to identify adjoining Sfs.

Still further, the present invention provides for a method of preparing a sample containing target nucleic acids by PCR, without purifying the PCR products prior to the SBH reactions. In Format I SBH, crude PCR products are applied to a substrate without  
10 prior purification, and the substrate may be washed prior to introduction of the labeled probes.

Still further, the present invention provides a method and an apparatus for analyzing a target nucleic acid. The apparatus comprises two arrays of nucleic acids that are mixed together at the desired time. In a preferred embodiment, the nucleic acids in  
15 one of the arrays are labeled. In a more preferred embodiment, a material is disposed between the two arrays and this material prevents the mixing of nucleic acids in the arrays. When this material is removed, or rendered permeable, the nucleic acids in the two arrays are mixed together. In an alternative preferred embodiment, the nucleic acids in one array are target nucleic acids and the nucleic acids in the other are oligonucleotide  
20 probes. In another preferred embodiment, the nucleic acids in both arrays are oligonucleotide probes. In another preferred embodiment, the nucleic acids in one array are oligonucleotide probes and target nucleic acids, and nucleic acids in the other array are oligonucleotide probes. In another preferred embodiment, the nucleic acids in both arrays are oligonucleotide probes and target nucleic acids.

25 One method of the present invention using the apparatus described above comprises the steps of providing an array of nucleic acids fixed to a substrate, providing a second array of nucleic acids, providing conditions that allow the nucleic acids in the second array to come into contact with the nucleic acids of the fixed array wherein one of the arrays of nucleic acids are target nucleic acids and the other array is oligonucleotide  
30 probes, and analyzing the hybridization results. In a preferred embodiment, the fixed array is target nucleic acid and the second array is labeled oligonucleotide probes. In a more preferred embodiment, there is a material disposed between the two arrays that

WO 99/09217

PCT/US98/16966

prevents mixing of the nucleic acids until the material is removed or rendered permeable to the nucleic acids.

In a second method of the present invention using the apparatus described above comprises the steps of providing two arrays of nucleic acid probes, providing conditions  
5 that allow the two arrays of probes to come into contact with each other and a target nucleic acid, ligating together probes that are adjacent on the target nucleic acid, and analyzing the results. In a preferred embodiment, the probes in one array are fixed and the probes in the other array are labeled. In a more preferred embodiment, there is a material disposed between the two arrays that prevents mixing of the probes until the  
10 material is removed or rendered permeable to the probes.

Still further, the present invention provides substrates on which arrays of oligonucleotide probes are fixed, wherein each probe is separated from its neighboring probes by a physical barrier that is resistant to the flow of the sample solution. In a preferred embodiment, the physical barrier is made of a hydrophobic material.

15 Still further, the present invention provides a method for making the arrays of oligonucleotide probes that are separated by physical barriers. In a preferred embodiment, a grid is applied to the substrate using an ink-jet head that applies a material which reduces the reaction volume of the array.

Still further, the present invention provides substrates on which oligonucleotides  
20 are fixed to form a three-dimensional array. The three-dimensional array combines high resolution for reading probe results (each level has a relatively low density of probes per  $\text{cm}^2$ ), with high information content in three dimensional space (multiple levels or probes).

Still further, the present invention provides a substrate to which oligonucleotide  
25 probes are fixed, wherein the oligonucleotide probes have spacers, and wherein the spacers increase the distance between the substrate and the informational portion of the oligonucleotide probe (e.g., the portion of the oligonucleotide probe which binds to the target and gives sequence information). In a preferred embodiment, the spacer comprises ribose sugars and phosphates, wherein the phosphates covalently bind the ribose sugars  
30 into a polymer by forming esters with the ribose sugars through their 5' and 3' hydroxyl groups.

WO 99/09217

PCT/US98/16966

Still further, the present invention provides a method for clustering cDNA clones into groups of similar or identical sequences, so that single representative clones may be selected from each group for sequencing. In a preferred embodiment, the method for clustering is used in the sequencing of a plurality of clones, comprising the steps of:

- 5 interrogating each clone with a plurality of oligonucleotide probes; determining which probes bind to each clone and the signal intensity for each probe; clustering clones into a plurality of groups by identifying clones that bind to similar probes with similar intensities; and sequencing at least one clone from each group. In a more preferred embodiment, the plurality of probes comprises from about 50 to about 500 different  
10 probes. In another more preferred embodiment, the plurality of probe comprises about 300 different probes. In a most preferred embodiment, the plurality of clones are a plurality of cDNA clones.

- Still further, the invention relates to oligonucleotide probes complexed (covalent or noncovalent) to discrete particles wherein the particles can be grouped into a plurality of  
15 sets based on a physical property. In a preferred embodiment, a different probe is attached to the discrete particles of each set, and the identity of the probe is determined by identifying the physical property of the discrete particles. In an alternative embodiment, the probe is identified on the basis of a physical property of the probe. The physical property includes any that can be used to differentiate the discrete particles, and  
20 includes, for example, size, fluorescence, radioactivity, electromagnetic charge, or absorbance, or label(s) may be attached to the particle such as a dye, a radionuclide, or an EML. In a preferred embodiment, discrete particles are separated by a flow cytometer which detects the size, charge, fluorescence, or absorbance of the particle.

- The invention also relates to methods using the probes complexed with the discrete  
25 particles to analyze target nucleic acids. These probes may be used in any of the methods described above, with the modification of identifying the probe by the physical property of the discrete particle. These probes may also be used in a format III approach where the "free" probe is identified by a label, and the probe complexed to the discrete particle is identified by the physical property. In a preferred embodiment, the probes are used to  
30 sequence a target nucleic acid using SBH.

The invention also relates to methods using agents which destabilize the binding of complementary polynucleotide strands (decrease the binding energy), or increase stability

WO 99/09217

PCT/US98/16966

of binding between complementary polynucleotide strands (increase the binding energy). In preferred embodiments, the agent is a trialkyl ammonium salt, sodium chloride, phosphate salts, borate salts, organic solvents such as formamide, glycol, dimethylsulfoxide, and dimethylformamide. urea, guanidinium, amino acid analogs such as betaine, polyamines such as spermidine and spermine, or other positively charged molecules which neutralize the negative charge of the phosphate backbone, detergents such as sodium dodecyl sulfate, and sodium lauryl sarcosinate, minor/major groove binding agents, positively charged polypeptides, and intercalating agents such as acridine, ethidium bromide, and anthracene. In a preferred embodiment, an agent is used to reduce or increase the  $T_m$  of a pair of complementary polynucleotides. In a more preferred embodiment, a mixture of the agents is used to reduce or increase the  $T_m$  of a pair of complementary polynucleotides. In a most preferred embodiment, an agent or a mixture of agents is used to increase the discrimination of perfect matches from mismatches for complementary polynucleotides. In a preferred embodiment, the agent or agents are added so that the binding energy from an AT base pair is approximately equivalent to the binding energy of a GC base pair. The energy of binding of these complementary polynucleotides may be increased by adding an agent that neutralizes or shields the negative charges of the phosphate groups in the polynucleotide backbone.

## 20 **BRIEF DESCRIPTION OF THE DRAWINGS**

Figure 1 illustrates a top view of the apparatus for mass producing probe arrays.

Figure 2 illustrates a side view of the apparatus for mass producing probe arrays.

Figure 3 is an exploded side view of the dispensing unit of the apparatus for mass producing probe arrays.

25

## **DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS**

Format 1 SBH is appropriate for the simultaneous analysis of a large set of samples. Parallel scoring of thousands of samples on large arrays may be performed in thousands of independent hybridization reactions using small pieces of membranes. The identification of DNA may involve 1-20 probes per reaction and the identification of mutations may in some cases involve more than 1000 probes specifically selected or designed for each sample. For identification of the nature of the mutated DNA segments,

WO 99/09217

PCT/US98/16966

specific probes may be synthesized or selected for each mutation detected in the first round of hybridizations.

DNA samples may be prepared in small arrays which may be separated by appropriate spacers, and which may be simultaneously tested with probes selected from a set of oligonucleotides which may be arrayed in multiwell plates. Small arrays may consist of one or more samples. DNA samples in each small array may include mutants or individual samples of a sequence. Consecutive small arrays may be organized into larger arrays. Such larger arrays may include replication of the same small array or may include arrays of samples of different DNA fragments. A universal set of probes includes sufficient probes to analyze a DNA fragment with prespecified precision, e.g. with respect to the redundancy of reading each base pair ("bp"). These sets may include more probes than are necessary for one specific fragment, but may include fewer probes than are necessary for testing thousands of DNA samples of different sequence.

DNA or allele identification and a diagnostic sequencing process may include the steps of:

- 1) Selection of a subset of probes from a dedicated, representative or universal set to be hybridized with each of a plurality of small arrays;
- 2) Adding a first probe to each subarray on each of the arrays to be analyzed in parallel;
- 3) Performing hybridization and scoring of the hybridization results;
- 4) Stripping off previously used probes;
- 5) Repeating hybridization, scoring and stripping steps for the remaining probes which are to be scored;
- 5) Processing the obtained results to obtain a final analysis or to determine additional probes to be hybridized;
- 6) Performing additional hybridizations for certain subarrays; and
- 7) Processing complete sets of data and obtaining a final analysis.

This approach provides fast identification and sequencing of a small number of nucleic acid samples of one type (e.g. DNA, RNA), and also provides parallel analysis of many sample types in the form of subarrays by using a presynthesized set of probes of manageable size. Two approaches have been combined to produce an efficient and



WO 99/09217

PCT/US98/16966

versatile process for the determination of DNA identity, for DNA diagnostics, and for identification of mutations.

For the identification of known sequences, a small set of shorter probes may be used in place of a longer unique probe. In this approach, although there may be more  
5 probes to be scored, a universal set of probes may be synthesized to cover any type of sequence. For example, a full set of 6-mers includes only 4,096 probes, and a complete set of 7-mers includes only 16,384 probes.

Full sequencing of a DNA fragment may be performed with two levels of hybridization. One level is hybridization of a sufficient set of probes that cover every base  
10 at least once. For this purpose, a specific set of probes may be synthesized for a standard sample. The results of hybridization with such a set of probes reveal whether and where mutations (differences) occur in non-standard samples. Further, this set of probes may include "negative" probes to confirm the hybridization results of the "positive" probes. To determine the identity of the changes, additional specific probes may be hybridized to  
15 the sample. This additional set of probes will have both "positive" (the mutant sequence) and "negative" probes, and the sequence changes will be identified by the positive probes and confirmed by the negative probes.

In another embodiment, all probes from a universal set may be scored. A universal set of probes allows scoring of a relatively small number of probes per sample  
20 in a two step process without an undesirable expenditure of time. The hybridization process may involve successive probings, in a first step of computing an optimal subset of probes to be hybridized first and, then, on the basis of the obtained results, a second step of determining additional probes to be scored from among those in a universal set. Both sets of probes have "negative" probes that confirm the positive probes in the set.  
25 Further, the sequence that is obtained may then be confirmed in a separate step by hybridizing the sample with a set of "negative" probes identified from the SBH results.

In SBH sequence assembly, K -1 oligonucleotides which occur repeatedly in analyzed DNA fragments due to chance or biological reasons may be subject to special consideration. If there is no additional information, relatively small fragments of DNA  
30 may be fully assembled in as much as every base pair is read several times.

In the assembly of relatively longer fragments, ambiguities may arise due to the repeated occurrence in a set of positively-scored probes of a K-1 sequence (i.e., a

WO 99/09217

PCT/US98/16966

sequence shorter than the length of the probe). This problem does not exist if mutated or similar sequences have to be determined (i.e., the K-1 sequence is not identically repeated). Knowledge of one sequence may be used as a template to correctly assemble a sequence known to be similar (e.g. by its presence in a database) by arraying the positive  
5 probes for the unknown sequence to display the best fit on the template.

The use of an array of sample avoids consecutive scoring of many oligonucleotides on a single sample or on a small set of samples. This approach allows the scoring of more probes in parallel by manipulation of only one physical object. Subarrays of DNA samples 1000 bp in length may be sequenced in a relatively short period of time. If the  
10 samples are spotted at 50 subarrays in an array and the array is reprobated 10 times, 500 probes may be scored. In screening for the occurrence of a mutation, enough probes may be used to cover each base three times. If a mutation is present, several covering probes will be affected. The use of information about the identity of negative probes may map the mutation with a two base precision. To solve a single base mutation mapped in this  
15 way, an additional 15 probes may be employed. These probes cover any base combination for two questionable positions (assuming that deletions and insertions are not involved). These probes may be scored in one cycle on 50 subarrays which contain a given sample. In the implementation of a multiple label color scheme (i.e., multiplexing), two to six probes, each having a different label such as a different fluorescent dye, may be used as a  
20 pool, thereby reducing the number of hybridization cycles and shortening the sequencing process.

In more complicated cases, there may be two close mutations or insertions. They may be handled with more probes. For example, a three base insertion may be solved with 64 probes. The most complicated cases may be approached by several steps of  
25 hybridization, and the selecting of a new set of probes on the basis of results of previous hybridizations.

If subarrays to be analyzed include tens or hundreds of samples of one type, then several of them may be found to contain one or more changes (mutations, insertions, or deletions). For each segment where mutation occurs, a specific set of probes may be  
30 scored. The total number of probes to be scored for a type of sample may be several hundreds. The scoring of replica arrays in parallel facilitates scoring of hundreds of probes in a relatively small number of cycles. In addition, compatible probes may be

WO 99/09217

PCT/US98/16966

pooled. Positive hybridizations may be assigned to the probes selected to check particular DNA segments because these segments usually differ in 75% of their constituent bases.

By using a larger set of longer probes, longer targets may be analyzed. These targets may represent pools of fragments such as pools of exon clones.

- 5 A specific hybridization scoring method may be employed to define the presence of mutants in a genomic segment to be sequenced from a diploid chromosomal set. Two variations are where: i) the sequence from one chromosome represents a known allele and the sequence from the other represents a new mutant; or, ii) both chromosomes contain new, but different mutants. In both cases, the scanning step designed to map changes  
10 gives a maximal signal difference of two-fold at the mutant position. Further, the method can be used to identify which alleles of a gene are carried by an individual and whether the individual is homozygous or heterozygous for that gene.

- Scoring two-fold signal differences required in the first case may be achieved efficiently by comparing corresponding signals with homozygous and heterozygous  
15 controls. This approach allows determination of a relative reduction in the hybridization signal for each particular probe in a given sample. This is significant because hybridization efficiency may vary more than two-fold for a particular probe hybridized with different nucleic acid fragments having the same full match target. In addition, different mutant sites may affect more than one probe depending upon the number of  
20 oligonucleotide probes. Decrease of the signal for two to four consecutive probes produces a more significant indication of a mutant site. Results may be checked by testing with small sets of selected probes among which one or few probes selected to give a full match signal which is on average eight-fold stronger than the signals coming from mismatch-containing duplexes.

- 25 Partitioned membranes allow a very flexible organization of experiments to accommodate relatively larger numbers of samples representing a given sequence type, or many different types of samples represented with relatively small numbers of samples. A range of 4-256 samples can be handled with particular efficiency. Subarrays within this range of numbers of dots may be designed to match the configuration and size of standard  
30 multiwell plates used for storing and labeling oligonucleotides. The size of the subarrays may be adjusted for different number of samples, or a few standard subarray sizes may be used. If all samples of a type do not fit in one subarray, additional subarrays or

WO 99/09217

PCT/US98/16966

membranes may be used and processed with the same probes. In addition, by adjusting the number of replicas for each subarray, the time for completion of identification or sequencing process may be varied.

As used herein, "intermediate fragment" means an oligonucleotide between 5 and 1000 bases in length, and preferably between 10 and 40 bp in length.

In Format 3, a first set of oligonucleotide probes of known sequence is immobilized on a solid support under conditions which permit them to hybridize with nucleic acids having respectively complementary sequences. A labeled, second set of oligonucleotide probes is provided in solution. Both within the sets and between the sets the probes may be of the same length or of different lengths. A nucleic acid to be sequenced or intermediate fragments thereof may be applied to the first set of probes in double-stranded form (especially where a recA protein is present to permit hybridization under non-denaturing conditions), or in single-stranded form and under conditions which permit hybrids of different degrees of complementarity (for example, under conditions which allow discrimination between full match and one base pair mismatch hybrids). The nucleic acid to be sequenced or intermediate fragments thereof may be applied to the first set of probes before, after or simultaneously with the second set of probes. Probes that bind to adjacent sites on the target are bound together (e.g., by stacking interactions or by a ligase or other means of causing chemical bond formation between the adjacent probes). After permitting adjacent probes to be bound, fragments and probes which are not immobilized to the surface by chemical bonding to a member of the first set of probe are washed away, for example, using a high temperature (up to 100 degrees C) wash solution which melts hybrids. The bound probes from the second set may then be detected using means appropriate to the label employed (which may, for example, be chemiluminescent, fluorescent, radioactive, enzymatic, densitometric, or electrophore mass labels).

Herein, nucleotide bases "match" or are "complementary" if they form a stable duplex by hydrogen bonding under specified conditions. For example, under conditions commonly employed in hybridization assays, adenine ("A") matches thymine ("T"), but not guanine ("G") or cytosine ("C"). Similarly, G matches C, but not A or T. Other bases which will hydrogen bond in less specific fashion, such as inosine or the Universal Base ("M" base, Nichols et al 1994), or other modified bases, such as methylated bases,

WO 99/09217

PCT/US98/16966

for example, are complementary to those bases for which they form a stable duplex under specified conditions. A probe is said to be "perfectly complementary" or is said to be a "perfect match" if each base in the probe forms a duplex by hydrogen bonding to a base in the nucleic acid to be sequenced according to the Watson and Crick base pairing rules

5 (i.e., absent any surrounding sequence effects, the duplex formed has the maximal binding energy for a particular probe). "Perfectly complementary" and "perfect match" are also meant to encompass probes which have analogs or modified nucleotides. A "perfect match" for an analog or modified nucleotide is judged according to a "perfect match rule" selected for that analog or modified nucleotide (e.g., the binding pair that has

10 maximal binding energy for a particular analog or modified nucleotide). Each base in a probe that does not form a binding pair according to the "rules" is said to be a "mismatch" under the specified hybridization conditions.

A list of probes may be assembled wherein each probe is a perfect match to the nucleic acid to be sequenced. The probes on this list may then be analyzed to order them

15 in maximal overlap fashion. Such ordering may be accomplished by comparing a first probe to each of the other probes on the list to determine which probe has a 3' end which has the longest sequence of bases identical to the sequence of bases at the 5' end of a second probe. The first and second probes may then be overlapped, and the process may be repeated by comparing the 5' end of the second probe to the 3' end of all of the

20 remaining probes and by comparing the 3' end of the first probe with the 5' end of all of the remaining probes. The process may be continued until there are no probes on the list which have not been overlapped with other probes. Alternatively, more than one probe may be selected from the list of positive probes, and more than one set of overlapped probes ("sequence nucleus") may be generated in parallel. The list of probes for either

25 such process of sequence assembly may be the list of all probes which are perfectly complementary to the nucleic acid to be sequenced or may be any subset thereof.

The 5' and 3' ends of the probes may be overlapped to generate longer stretches of sequence. This process of assembling probes continues until an ambiguity arises because of a branch point (a probe is repeated in the fragment), repetitive sequences

30 longer than the probes, or an uncloned segment. The stretches of sequence between any two ambiguities are referred to as fragment of a subclone sequence (Sfs). Where ambiguities arise in sequence assembly due to the availability of alternative proper

WO 99/09217

PCT/US98/16966

overlaps with probes, hybridization with longer probes spanning the site of overlap alternatives, competitive hybridization, ligation of alternative end to end pairs of probes spanning the site of ambiguity or single pass gel analysis (to provide an unambiguous ordering of Sfs) may be used.

- 5 By employing the above procedures, one may obtain any desired level of sequence, from a pattern of hybridization (which may be correlated with the identity of a nucleic acid sample to serve as a signature for identifying the nucleic acid sample) to overlapping or non-overlapping probes up through assembled Sfs and on to complete sequence for an intermediate fragment or an entire source DNA molecule (e.g. a  
10 chromosome).

Sequencing may generally comprise the following steps:

- (a) contacting an array of immobilized oligonucleotide probes with a nucleic acid fragment under conditions effective to allow the fragment to form a primary complex with an immobilized probe having a complementary sequence;
- 15 (b) contacting this primary complex with a set of labeled oligonucleotide probes in solution under conditions effective to allow the primary complex to hybridize to the labeled probe, thereby forming secondary complexes wherein the fragment is hybridized with both an immobilized probe and a labeled probe;
- (c) removing from a secondary complex any labeled probe that has not  
20 hybridized adjacent to an immobilized probe;
- (d) detecting the presence of adjacent labeled and unlabeled probes by detecting the presence of the label; and
- (e) determining a nucleotide sequence of the fragment by connecting the known sequence of the immobilized and labeled probes.

- 25 Hybridization and washing conditions may be selected to detect substantially perfect match hybrids (such as those wherein the fragment and probe hybridize at six out of seven positions), may be selected to allow differentiation of perfect matches and one base pair mismatches, or may be selected to permit detection only of perfect match hybrids.

- 30 Suitable hybridization conditions may be routinely determined by optimization procedures or pilot studies. Such procedures and studies are routinely conducted by those skilled in the art to establish protocols for use in a laboratory. See e.g., Ausubel et al.,

WO 99/09217

PCT/US98/16966

*Current Protocols in Molecular Biology*, Vol. 1-2, John Wiley & Sons (1989); Sambrook et al., *Molecular Cloning A Laboratory Manual*, 2nd Ed., Vols. 1-3, Cold Springs Harbor Press (1989); and Maniatis et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Cold Spring Harbor, New York (1982), all of which are  
5 incorporated by reference herein. For example, conditions such as temperature, concentration of components, hybridization and washing times, buffer components, and their pH and ionic strength may be varied.

In embodiments wherein the labeled and immobilized probes are not physically or chemically linked, detection may rely solely on washing steps of controlled stringency.  
10 Under such conditions, adjacent probes have increased binding affinity because of stacking interactions between the adjacent probes. Conditions may be varied to optimize the process as described above.

In embodiments wherein the immobilized and labeled probes are ligated, ligation may be implemented by a chemical ligating agent (e.g. water-soluble carbodiimide or  
15 cyanogen bromide), or a ligase enzyme, such as the commercially available T<sub>4</sub> DNA ligase may be employed. The washing conditions may be selected to distinguish between adjacent versus nonadjacent labeled and immobilized probes exploiting the difference in stability for adjacent probes versus nonadjacent probes.

Oligonucleotide probes may be labeled with fluorescent dyes, chemiluminescent  
20 systems, radioactive labels (e.g., <sup>35</sup>S, <sup>3</sup>H, <sup>32</sup>P or <sup>33</sup>P) or with isotopes detectable by mass spectrometry.

Where a nucleic acid molecule of unknown sequence is longer than about 45 or 50 bp, the molecule may be fragmented and the sequences of the fragments determined. Fragmentation may be accomplished by restriction enzyme digestion, shearing or NaOH.  
25 Fragments may be separated by size (e.g. by gel electrophoresis) to obtain a preferred fragment length of about ten to forty bps.

Oligonucleotides may be immobilized, by a number of methods known to those skilled in the art, such as laser-activated photodeprotection attachment through a phosphate group using reagents such as a nucleoside phosphoramidite or a nucleoside  
30 hydrogen phosphorate. Glass, nylon, silicon and fluorocarbon supports may be used.

Oligonucleotides may be organized into arrays, and these arrays may include all or a subset of all probes of a given length, or sets of probes of selected lengths.

WO 99/09217

PCT/US98/16966

Hydrophobic partitions may be used to separate probes or subarrays of probes. Arrays may be designed for various applications (e.g. mapping, partial sequencing, sequencing of targeted regions for diagnostic purposes, mRNA sequencing and large scale sequencing). A specific chip may be designed to be dedicated to a particular application by selecting a  
5 combination and arrangement of probes on a substrate.

For example, 1024 immobilized probe arrays of all oligonucleotide probes 5 bases in length (each array containing 1024 distinct probes) may be constructed. The probes in this example are 5-mers in an informational sense (they may actually be longer probes). A second set of 1024 5-mer probes may be labeled, and one of each labeled probe may be  
10 applied to an array of immobilized probes along with a fragment to be sequenced. In this example, 1024 arrays would be combined in a large superarray, or "superchip." In those instances where an immobilized probe and one of the labeled probes hybridize end-to-end along a nucleic acid fragment., the two probes are joined. for example by ligation, and, after removing unbound label, 10-mers complementary to the sample  
15 fragment are detected by the correlation of the presence of a label at a point in an array having an immobilized probe of known sequence to which was applied a labeled probe of known sequence. The sequence of the sample fragment is simply the sequence of the immobilized probe continued in the sequence of the labeled probe. In this way, all one million possible 10-mers may be tested by a combinatorial process which employs only  
20 5-mers and which thus involves one thousandth of the amount of effort for oligonucleotide synthesis.

In a preferred embodiment, the substrate which supports the array of oligonucleotide probes is partitioned into sections so that each probe in the array is separated from adjacent probes by a physical barrier which may be, for example, a  
25 hydrophobic material. In a preferred embodiment, the physical barrier has a width of from 100  $\mu\text{m}$  to 30  $\mu\text{m}$ . In a more preferred embodiment, the distance from the center of each probe to the center of any adjacent probes is 325  $\mu\text{m}$ . These arrays of probes may be "mass-produced" using a nonmoving, fixed substrate or a substrate fixed to a rotating drum or plate with an ink-jet deposition apparatus, for example, a microdrop dosing head;  
30 and a suitable robotic system, for example, an anorad gantry.

In an alternative preferred embodiment, the oligonucleotide probes are fixed to a three-dimensional array. The three-dimensional array is comprised of multiple layers,



WO 99/09217

PCT/US98/16966

- and each layer may be analyzed separate and apart from the other layers. The three dimensional array may take a number of forms, including, for example, the array may be disposed on a substrate having multiple depressions with probes located at different depths within the depressions (each level is made up of probes at similar depths within the depression); or the array may be disposed on a substrate having depressions of different depths with the probes located at the bottom of the depression, or at the peaks separating the depressions or some combination of peaks and depressions may be used (each level is made up of all the probes at a certain depth); or the array may be disposed on a substrate comprised of multiple sheets that are layered to form a three-dimensional array.
- 10 The probes in these arrays may include spacers that increase the distance between the surface of the substrate and the informational portion of the probes. The spacers may be comprised of atoms capable of forming at least two covalent bonds such as carbon, silicon, oxygen, sulfur, phosphorous, and the like, or may be comprised of molecules capable of forming at least two covalent bonds such as sugar-phosphate groups, amino acids, peptides, nucleosides, nucleotides, sugars, carbohydrates, aromatic rings, hydrocarbon rings, linear and branched hydrocarbons, and the like.

- A nucleic acid sample to be sequenced may be fragmented or otherwise treated (for example, by the use of *recA*) to avoid hindrance to hybridization from secondary structure in the sample. The sample may be fragmented by, for example, digestion with a restriction enzyme such as *Cvi* II, physical shearing (e.g. by ultrasound), or by NaOH treatment. The resulting fragments may be separated by gel electrophoresis and fragments of an appropriate length, such as between about 10 bp and about 40 bp, may be extracted from the gel. In a preferred embodiment, the "fragments" of the nucleic acid sample cannot be ligated to other fragments in the pool. Such a pool of fragments may be obtained by treating the fragmented nucleic acids with a phosphatase (e.g., calf intestinal phosphatase). Alternatively, nonligatable fragments of the sample nucleic acid may be obtained by using random primers (e.g.,  $N_5-N_9$ , where  $N = A, G, T, \text{ or } C$ ) in a Sanger-dideoxy sequencing reaction with the sample nucleic acid. This will produce fragments of DNA that have a complementary sequence to the target nucleic acid and that are terminated in a dideoxy residue that cannot be ligated to other fragments.

A reusable Format 3 SBH array may be produced by introducing a cleavable bond between the fixed and labeled probes and then cleaving this bond after a round of Format

WO 99/09217

PCT/US98/16966

3 analyzes is finished. The labeled probes may be ribonucleotides or a ribonucleotide may be used as the joining base in the labeled probe so that this probe may subsequently be removed, e.g., by RNase or uracil-DNA glycosylate treatment, or NaOH treatment. In addition, bonds produced by chemical ligation may be selectively cleaved.

5 Other variations include the use of modified oligonucleotides to increase specificity or efficiency, cycling hybridizations to increase the hybridization signal, for example by performing a hybridization cycle under conditions (e.g. temperature) optimally selected for a first set of labeled probes followed by hybridization under conditions optimally selected for a second set of labeled probes. Shifts in reading frame may be determined  
10 by using mixtures (preferably mixtures of equimolar amounts) of probes ending in each of the four nucleotide bases A, T, C and G.

Branch points produce ambiguities as to the ordered sequence of a fragment. Although the sequence information is determined by SBH, either: (i) long read length, single-pass gel sequencing at a fraction of the cost of complete gel sequencing; or (ii)  
15 comparison to related sequences, may be used to order hybridization data where such ambiguities ("branch points") occur. Primers for single pass gel sequencing through the branch points are identified from the SBH sequence information or from known vector sequences, e.g., the flanking sequences to the vector insert site, and standard Sanger-sequencing reactions are performed on the sample nucleic acid. The sequence obtained  
20 from this single pass gel sequencing is compared to the Sfs that read into and out of the branch points to identify the order of the Sfs. Alternatively, the Sfs may be ordered by comparing the sequence of the Sfs to related sequences and ordering the Sfs to produce a sequence that is closest to the related sequence.

In addition, the number of tandem repetitive nucleic acid segments in a target  
25 fragment may be determined by single-pass gel sequencing. As tandem repeats occur rarely in protein-encoding portions of a gene, the gel-sequencing step will be performed only when one of these noncoding regions is identified as being of particular interest (e.g., if it is an important regulatory region).

Obtaining information about the degree of hybridization exhibited for a set of only  
30 about 200 oligonucleotides probes (about 5% of the effort required for complete sequencing) defines a unique signature of each gene and may be used for sorting the cDNAs from a library to determine if the library contains multiple copies of the same

WO 99/09217

PCT/US98/16966

gene. By such signatures, identical, similar and different cDNAs can be distinguished and inventoried.

Nucleic acids and methods for isolating, cloning and sequencing nucleic acids are well known to those of skill in the art. See e.g., Ausubel et al., *Current Protocols in Molecular Biology*, Vol. 1-2, John Wiley & Sons (1989); and Sambrook et al., *Molecular Cloning A Laboratory Manual*, 2nd Ed., Vols. 1-3, Cold Springs Harbor Press (1989), both of which are incorporated by reference herein.

SBH is a well developed technology that may be practiced by a number of methods known to those skilled in the art. Specifically, techniques related to sequencing by hybridization of the following documents is incorporated by reference herein:

Drmanac et al., U.S. Patent No. 5,202,231 (hereby incorporated by reference herein) - Issued April 13, 1993; Drmanac et al., *Genomics*, 4, 114-128 (1989); Drmanac et al., *Proceedings of the First Int'l. Conf. Electrophoresis Supercomputing Human Genome* Cantor et al. eds, World Scientific Pub. Co., Singapore, 47-59 (1991); Drmanac et al., *Science*, 260, 1649-1652 (1993); Lehrach et al., *Genome Analysis: Genetic and Physical Mapping*, 1, 39-81 (1990), Cold Spring Harbor Laboratory Press; Drmanac et al., *Nucl. Acids Res.*, 4691 (1986); Stevanovic et al., *Gene*, 79, 139 (1989); Panusku et al., *Mol. Biol. Evol.*, 1, 607 (1990); Nizetic et al., *Nucl. Acids Res.*, 19, 182 (1991); Drmanac et al., *J. Biomol. Struct. Dyn.*, 5, 1085 (1991); Hoheisel et al., *Mol. Gen.*, 4, 125-132 (1991); Strezoska et al., *Proc. Nat'l. Acad. Sci. (USA)*, 88, 10089 (1991); Drmanac et al., *Nucl. Acids Res.*, 19, 5839 (1991); and Drmanac et al., *Int. J. Genome Res.*, 1, 59-79 (1992).

The present invention is illustrated in the following examples. Upon consideration of the present disclosure, one of skill in the art will appreciate that many other embodiments and variations may be made in the scope of the present invention. Accordingly, it is intended that the broader aspects of the present invention not be limited to the disclosure of the following examples.

30

WO 99/09217

PCT/US98/16966

**EXAMPLE 1****Preparation of Sets of Probes**

Two types of universal sets of probes may be prepared. The first is a complete set (or at least a noncomplementary subset) of relatively short probes, for example all 4096  
5 (or about 2000 non-complementary) 6-mers, or all 16,384 (or about 8,000 non-complementary) 7-mers. Full noncomplementary subsets of 8-mers and longer probes are less convenient inasmuch as they include 32,000 or more probes.

A second type of probe set is selected as a small subset of probes still sufficient for reading every bp in any sequence with at least with one probe. For example, 12 of 16  
10 dimers are sufficient. A small subset for 7-mers, 8-mer and 9-mers for sequencing double stranded DNA may be about 3000, 10,000 and 30,000 probes, respectively.

Sets of probes may also be selected to identify a target nucleic acid of known sequence, and/or to identify alleles or mutants of a target nucleic acid with a known sequence. Such a set of probes contains sufficient probes so that every nucleotide  
15 position of the target nucleic acid is read at least once. Alleles or mutants are identified by the loss of binding of one of the "positive" probes. The specific sequence of these alleles or mutants is then determined by interrogating the target nucleic acid with sets of probes that contain every possible nucleotide change and combination of changes at these probe positions.

20 Sets of probes may also be comprised of from 50 probes to a universal set of probes (all probes of a certain length), more preferably the set is comprised of 100-500 probes, and in a most preferred embodiment, the probe set contains 300 probes. In a preferred embodiment, the set of probes are 6-9 nucleotides in length, and are used to cluster cDNA clones into groups of similar or identical sequences, so that single  
25 representative clones may be selected from each group for sequencing.

Probes may be prepared using standard chemistry with one to three non-specified (mixed A,T,C and G) or universal (e.g. M base or inosine) bases at the ends. If radiolabelling is used, probes may have an OH group at the 5' end for kinasing by radiolabelled phosphorous groups. Alternatively, probes labelled with any compatible  
30 system, such as fluorescent dyes, may be employed. Other types of probes, such as PNA (Protein Nucleic Acids) or probes containing modified bases which change duplex stability also may be used.

WO 99/09217

PCT/US98/16966

Probes may be stored in bar-coded multiwell plates. For small numbers of probes, 96-well plates may be used; for 10,000 or more probes, storage in 384- or 864-well plates is preferred. Stacks of 5 to 50 plates are enough to store all probes. Approximately 5 pg of a probe may be sufficient for hybridization with one DNA sample. Thus, from a small synthesis of about 50 mg per probe, ten million samples may be analyzed. If each probe is used for every third sample, and if each sample is 1000 bp in length, then over 30 billion bases (10 human genomes) may be sequenced by a set of 5,000 probes.

## **EXAMPLE 2**

### **10 Probes Having Modified Oligonucleotides**

Modified oligonucleotides may be introduced into hybridization probes and used under appropriate conditions therefor. For example, pyrimidines with a halogen at the C<sup>5</sup>-position may be used to improve duplex stability by influencing base stacking. 2,6-diaminopurine may be used to provide a third hydrogen bond in base pairing with thymine, thereby thermally stabilizing DNA-duplexes. Using 2,5-diaminopurine may increase duplex stability to allow more stringent conditions for annealing, thereby improving the specificity of duplex formation, suppressing background problems and permitting the use of shorter oligomers.

The synthesis of the triphosphate versions of these modified nucleotides is disclosed by Hoheisel & Lehrach (1990).

One may also use the non-discriminatory base analogue, or universal base, as designed by Nichols et al. (1994). This new analogue, 1-(2 -deoxy-  
-D-ribofuranosyl)-3-nitropyrrole (designated M), was generated for use in oligonucleotide probes and primers for solving the design problems that arise as a result of the degeneracy of the genetic code, or when only fragmentary peptide sequence data are available. This analogue maximizes stacking while minimizing hydrogen-bonding interactions without sterically disrupting a DNA duplex.

The M nucleoside analogue was designed to maximize stacking interactions using aprotic polar substituents linked to heteroaromatic rings, enhancing intra- and inter-strand stacking interactions to lessen the role of hydrogen bonding in base-pairing specificity. Nichols *et al.* (1994) favored 3-nitropyrrole 2 -deoxyribonucleoside because of its

WO 99/09217

PCT/US98/16966

structural and electronic resemblance to p-nitroaniline, whose derivatives are among the smallest known intercalators of double-stranded DNA.

The dimethoxytrityl-protected phosphoramidite of nucleoside M is also available for incorporation into nucleotides used as primers for sequencing and polymerase chain reaction (PCR). Nichols *et al.* (1994) showed that a substantial number of nucleotides can be replaced by M without loss of primer specificity.

A unique property of M is its ability to replace long strings of contiguous nucleosides and still yield functional sequencing primers. Sequences with three, six and nine M substitutions have all been reported to give readable sequencing ladders, and PCR with three different M-containing primers all resulted in amplification of the correct product (Nichols *et al.*, 1994).

The ability of 3-nitropyrrole-containing oligonucleotides to function as primers strongly suggests that a duplex structure must form with complementary strands. Optical thermal profiles obtained for the oligonucleotide pairs d(5 -C<sub>2</sub>-T<sub>5</sub>XT<sub>5</sub>G<sub>2</sub>-3 ) and d(5 -C<sub>2</sub>A<sub>5</sub>YA<sub>5</sub>G<sub>2</sub>-3 ) (where X and Y can be A, C, G, T or M) were reported to fit the normal sigmoidal pattern observed for the DNA double-to single strand transition. The T<sub>m</sub> values of the oligonucleotides containing X M base pairs (where X was A, C, G or T, and Y was M) were reported to all fall within a 3°C range (Nichols *et al.*, 1994).

### 20 EXAMPLE 3

#### Selection and Labeling of Probes

When an array of subarrays is produced, the sets of probes to be hybridized in each of the hybridization cycles on each of the subarrays is defined. For example, a set of 384 probes may be selected from the universal set, and 96 probings may be performed in each of 4 cycles. Probes selected to be hybridized in one cycle preferably have similar G+C contents.

Selected probes for each cycle are transferred to a 96-well plate and then are labelled by kinasing or by other labeling procedures if they are not labelled (e.g. with stable fluorescent dyes) before they are stored.

30 On the basis of the first round of hybridizations, a new set of probes may be defined for each of the subarrays for additional cycles. Some of the arrays may not be used in some of the cycles. For example, if only 8 of 64 patient samples exhibit a

WO 99/09217

PCT/US98/16966

mutation and 8 probes are scored first for each mutation. then all 64 probes may be scored in one cycle and 32 subarrays are not used. These unused subarrays may then be treated with hybridization buffer to prevent drying of the filters.

Probes may be retrieved from the storing plates by any convenient approach, such as a single channel pipetting device, or a robotic station, such as a Beckman Biomek 1000 (Beckman Instruments, Fullerton, California) or a Mega Two robot (Megamation, Lawrenceville, New Jersey). A robotic station may be integrated with data analysis programs and probe managing programs. Outputs of these programs may be inputs for one or more robotic stations.

Probes may be retrieved one by one and added to subarrays covered by hybridization buffer. It is preferred that retrieved probes be placed in a new plate and labelled or mixed with hybridization buffer. The preferred method of retrieval is by accessing stored plates one by one and pipetting (or transferring by metal pins) a sufficient amount of each selected probe from each plate to specific wells in an intermediary plate. An array of individually addressable pipettes or pins may be used to speed up the retrieval process.

#### **EXAMPLE 4**

##### **Preparation of Labeled Probes**

The oligonucleotide probes may be prepared by automated synthesis, which is routine to those of skill in the art, for example, using an Applied Biosystems system. Alternatively, probes may be prepared using Genosys Biotechnologies Inc. Methods using stacks of porous Teflon wafers.

Oligonucleotide probes may be labeled with, for example, radioactive labels ( $^{35}\text{S}$ ,  $^{32}\text{P}$ ,  $^{33}\text{P}$ , and preferably,  $^{33}\text{P}$ ) for arrays with 100-200  $\mu\text{m}$  or 100-400  $\mu\text{m}$  spots; non-radioactive isotopes (Jacobsen *et al.*, 1990); or fluorophores (Brumbaugh *et al.*, 1988). All such labeling methods are routine in the art, as exemplified by the relevant sections in Sambrook *et al.* (1989) and by further references such as Schubert *et al.* (1990), Murakami *et al.* (1991) and Cate *et al.* (1991), all articles being specifically incorporated herein by reference.

WO 99/09217

PCT/US98/16966

In regard to radiolabelling, the common methods are end-labeling using T4 polynucleotide kinase or high specific activity labeling using Klenow or even T7 polymerase. These are described as follows.

Synthetic oligonucleotides are synthesized without a phosphate group at their 5  
5 termini and are therefore easily labeled by transfer of the  $^{32}\text{P}$  or  $^{33}\text{P}$  from [  $^{32}\text{P}$ ]ATP or [  $^{33}\text{P}$ ]ATP using the enzyme bacteriophage T4 polynucleotide kinase. If the reaction is carried out efficiently, the specificity activity of such probes can be as high as the specific activity of the [  $^{32}\text{P}$ ]ATP or [  $^{33}\text{P}$ ]ATP itself. The reaction described below is designed to label 10 pmoles of an oligonucleotide to high specific activity. Labeling of different  
10 amounts of oligonucleotide can easily be achieved by increasing or decreasing the size of the reaction, keeping the concentrations of all components constant.

A reaction mixture would be created using 1.0 ul of oligonucleotide (10 pmoles/ul); 2.0 ul of 10 x bacteriophage T4 polynucleotide kinase buffer; 5.0 ul of [  $^{32}\text{P}$ ]ATP or [  $^{33}\text{P}$ ]ATP (sp. Act. 5000 Ci/mmmole; 10 mCi/ml in aqueous solution) (10  
15 pmoles); and 11.4 ul of water. Eight (8) units (~ 1 ul) of bacteriophage T4 polynucleotide kinase is added to the reaction mixture, and incubated for 45 minutes at 37°C. The reaction is heated for 10 minutes at 68°C to inactivate the bacteriophage T4 polynucleotide kinase.

The efficiency of transfer of  $^{32}\text{P}$  or  $^{33}\text{P}$  to the oligonucleotide and its specific  
20 activity is then determined. If the specific activity of the probe is acceptable, it is purified. If the specific activity is too low, an additional 8 units of enzyme is added and incubated for a further 30 minutes at 37°C before heating the reaction for 10 minutes at 68°C to inactivate the enzyme.

Purification of radiolabeled oligonucleotides can be achieved by, e.g., precipitation  
25 with ethanol; precipitation with cetylpyridinium bromide; by chromatography through bio-gel P-60; or by chromatography on a Sep-Pak C<sub>18</sub> column, or by polyacrylamide gel electrophoresis.

Probes of higher specific activities can be obtained using the Klenow fragment of *E. coli*. DNA polymerase I to synthesize a strand of DNA complementary to the  
30 synthetic oligonucleotide. A short primer is hybridized to an oligonucleotide template whose sequence is the complement of the desired radiolabeled probe. The primer is then extended using the Klenow fragment of *E. coli* DNA polymerase I to incorporate [  $^{32}\text{P}$ ]



WO 99/09217

PCT/US98/16966

dNTPs or [ $^{33}\text{P}$ ] dNTPs in a template-directed manner. After the reaction, the template and product are separated by denaturation followed by electrophoresis through a polyacrylamide gel under denaturing conditions. With this method, it is possible to generate oligonucleotide probes that contain several radioactive atoms per molecule of

5 oligonucleotide.

To use this method, one would mix in a microfuge tube the calculated amounts of [ $\alpha$ - $^{32}\text{P}$ ]dNTPs or [ $\alpha$ - $^{33}\text{P}$ ]dNTPs necessary to achieve the desired specific activity and sufficient to allow complete synthesis of all template strands. Then add to the tube the appropriate amounts of primer and template DNAs, with the primer being in three- to

10 tenfold molar excess over the template.

0.1 volume of 10 x Klenow buffer would then be added and mixed well. 2-4 units of the Klenow fragment of *E. coli* DNA polymerase I would then be added per 5 ul of reaction volume, mixed and incubated for 2-3 hours at 4°C. If desired, the process of the reaction may be monitored by removing small (0.1 ul) aliquots and measuring the

15 proportion of radioactivity that has become precipitable with 10% trichloroacetic acid (TCA).

The reaction would be diluted with an equal volume of gel-loading buffer, heated to 80°C for 3 minutes, and then the entire sample loaded on a denaturing polyacrylamide gel. Following electrophoresis, the gel is autoradiographed, allowing the probe to be

20 localized and removed from the gel. Various methods for fluorescent probe labeling are also available, e.g., Brumbaugh *et al.* (1988) describe the synthesis of fluorescently labeled primers. A deoxyuridine analog with a primary amine "linker arm" of 12 atoms attached at C-5 is synthesized. Synthesis of the analog consists of derivatizing 2

25 -deoxyuridine through organometallic intermediates to give 5 (methyl propenoyl)-2 -deoxyuridine. Reaction with dimethoxytrityl-chloride produces the corresponding 5

-dimethoxytrityl adduct. The methyl ester is hydrolyzed, activated, and reacted with an appropriately monoacylated alkyl diamine. After purification, the resultant linker arm nucleosides are converted to nucleoside analogs suitable for chemical oligonucleotide synthesis.

Oligonucleotides would then be made that include one or two linker arm bases by

30 using modified phosphoridite chemistry. To a solution of 50 nmol of the linker arm oligonucleotide in 25 ul of 500 mM sodium bicarbonate (pH 9.4) is added 20 ul of 300

WO 99/09217

PCT/US98/16966

mM FITC in dimethyl sulfoxide. The mixture is agitated at room temperature for 6 hrs. The oligonucleotide is separated from free FITC by elution from a 1 x 30 cm Sephadex G-25 column with 20 mM ammonium acetate (pH 6), combining fractions in the first UV-absorbing peak.

- 5 In general, fluorescent labeling of an oligonucleotide at its 5'-end initially involved two steps. First, a *N*-protected aminoalkyl phosphoramidite derivative is added to the 5'-end of an oligonucleotide during automated nucleic acid synthesis. After removal of all protecting groups, the NHS ester of an appropriate fluorescent dye is coupled to the 5'-amino group overnight followed by purification of the labeled oligonucleotide from the  
10 excess of dye using reverse phase HPLC or PAGE.

Schubert *et al.* (1990) described the synthesis of a phosphoramidite that enables oligonucleotides labeled with fluorescein to be produced during automated DNA synthesis.

- Murakami *et al.* also described the preparation of fluorescein-labeled  
15 oligonucleotides.

Cate *et al.* (1991) describe the use of oligonucleotide probes directly conjugated to alkaline phosphatase in combination with a direct chemiluminescent substrate (AMPPD) to allow probe detection.

- Labeled probes could readily be purchased from a variety of commercial sources,  
20 including GENSET, rather than synthesized.

- Other labels include ligands which can serve as specific binding members to a labeled antibody, chemiluminescers, enzymes, antibodies which can serve as a specific binding pair member for a labeled ligand, and the like. A wide variety of labels have been employed in immunoassays which can readily be employed. Still other labels  
25 include antigens, groups with specific reactivity, and electrochemically detectable moieties.

In general, labeling of nucleic acids with electrophore mass labels ("EML") is described, for example, in Xu *et al.*, J. Chromatography 764:95-102 (1997).

- Electrophores are compounds that can be detected with high sensitivity by electron  
30 capture mass spectrometry (EC-MS). EMLs can be attached to a probe using chemistry that is well known in the art for reversibly modifying a nucleotide (e.g., well known nucleotide synthesis chemistry teaches a variety of methods for attaching molecules to

WO 99/09217

PCT/US98/16966

- nucleotides as protecting groups). EMLs are detected using a variety of well known electron capture mass spectrometry devices (e.g., devices sold by Finnigan Corporation). Further, techniques that may be used in the detection of EMLs include, for example, fast atomic bombardment mass spectrometry (see, e.g., Koster *et al.*, Biomedical Environ. Mass Spec. 14:111-116 (1987)); plasma desorption mass spectrometry; electrospray/ion spray (see, e.g., Fenn *et al.*, J. Phys. Chem. 88:4451-59 (1984), PCT Appln. No. WO 90/14148, Smith *et al.*, Anal. Chem. 62:882-89 (1990)); and matrix-assisted laser desorption/ionization (Hillenkamp, *et al.*, "Matrix Assisted UV-Laser Desorption/Ionization: A New Approach to Mass Spectrometry of Large Biomolecules," *Biological Mass Spectrometry* (Burlingame and McCloskey, eds.), Elsevier Science Publishers, Amsterdam, pp. 49-60, 1990); Huth-Fehre *et al.*, "Matrix Assisted Laser Desorption Mass Spectrometry of Oligodeoxythymidylic Acids," *Rapid Communications in Mass Spectrometry*, 6:209-13 (1992)).

- In preferred embodiments, the EMLs are attached to a probe by a covalent bond that is light sensitive. The EML is released from the probe after hybridization with a target nucleic acid by a laser or other light source emitting the desired wavelength of light. The EML is then fed into a GC-MS (gas chromatograph -mass spectrometer) or other appropriate device, and identified by its mass.

## 20 **EXAMPLE 5**

### **Preparation of Sequencing Chips and Arrays**

- A basic example is using 6-mers attached to 50 micron surfaces to give a chip with dimensions of 3 x 3 mm which can be combined to give an array of 20 x 20 cm. Another example is using 9-mer oligonucleotides attached to 10 x 10 microns surface to create a 9-mer chip, with dimensions of 5 x 5 mm. 4000 units of such chips may be used to create a 30 x 30 cm array. In an array in which 4,000 to 16,000 oligochips are arranged into a square array. A plate, or collection of tubes, as also depicted, may be packaged with the array as part of the sequencing kit.

- The arrays may be separated physically from each other or by hydrophobic surfaces. One possible way to utilize the hydrophobic strip separation is to use technology such as the Iso-Grid Microbiology System produced by QA Laboratories, Toronto, Canada.

WO 99/09217

PCT/US98/16966

Hydrophobic grid membrane filters (HGMF) have been in use in analytical food microbiology for about a decade where they exhibit unique attractions of extended numerical range and automated counting of colonies. One commercially-available grid is ISO-GRID™ from QA Laboratories Ltd. (Toronto, Canada) which consists of a square 5 (60 x 60 cm) of polysulfone polymer (Gelman Tuffryn HT-450, 0.45u pore size) on which is printed a black hydrophobic ink grid consisting of 1600 (40 x 40) square cells. HGMF have previously been inoculated with bacterial suspensions by vacuum filtration and incubated on the differential or selective media of choice.

Because the microbial growth is confined to grid cells of known position and size 10 on the membrane, the HGMF functions more like an MPN apparatus than a conventional plate or membrane filter. Peterkin *et al.* (1987) reported that these HGMFs can be used to propagate and store genomic libraries when used with a HGMF replicator. One such instrument replicates growth from each of the 1600 cells of the ISO-GRID and enables many copies of the master HGMF to be made (Peterkin *et al.*, 1987).

15 Sharpe *et al.* (1989) also used ISO-GRID HGMF from QA Laboratories and an automated HGMF counter (MI-100 Interpreter) and RP-100 Replicator. They reported a technique for maintaining and screening many microbial cultures.

Peterkin and colleagues later described a method for screening DNA probes using the hydrophobic grid-membrane filter (Peterkin *et al.*, 1989). These authors reported 20 methods for effective colony hybridization directly on HGMFs. Previously, poor results had been obtained due to the low DNA binding capacity of the epoxysulfone polymer on which the HGMFs are printed. However, Peterkin *et al.* (1989) reported that the binding of DNA to the surface of the membrane was improved by treating the replicated and incubated HGMF with polyethyleneimine, a polycation, prior to contact with DNA. 25 Although this early work uses cellular DNA attachment, and has a different objective to the present invention, the methodology described may be readily adapted for Format 3 SBH.

In order to identify useful sequences rapidly, Peterkin *et al.* (1989) used radiolabeled plasmid DNA from various clones and tested its specificity against the DNA 30 on the prepared HGMFs. In this way, DNA from recombinant plasmids was rapidly screened by colony hybridization against 100 organisms on HGMF replicates which can be easily and reproducibly prepared.

WO 99/09217

PCT/US98/16966

Manipulation with small (2-3 mm) chips, and parallel execution of thousands of the reactions. The solution of the invention is to keep the chips and the probes in the corresponding arrays. In one example, chips containing 250,000 9-mers are synthesized on a silicon wafer in the form of 8 x 8 mM plates (15 uM/oligonucleotide. Pease et al.,  
5 1994) arrayed in 8 x 12 format (96 chips) with a 1 mM groove in between. Probes are added either by multichannel pipette or pin array, one probe on one chip. To score all 4000 6-mers, 42 chip arrays have to be used, either using different ones. or by reusing one set of chip arrays several times.

In the above case, using the earlier nomenclature of the application.  $F=9$ ;  $P=6$ ;  
10 and  $F + P = 15$ . Chips may have probes of formula  $B_xN_n$ , where  $x$  is a number of specified bases  $B$ ; and  $n$  is a number of non-specified bases, so that  $x = 4$  to 10 and  $n = 1$  to 4. To achieve more efficient hybridization, and to avoid potential influence of any support oligonucleotides, the specified bases can be surrounded by unspecified bases, thus represented by a formula such as  $(N)_nB_x(N)_m$ .

15 In another embodiment of the chips, the substrate which supports the array of oligonucleotide probes is partitioned into sections so that each probe in the array is separated from adjacent probes by a physical barrier which may be, for example, a hydrophobic material. In a preferred embodiment, the physical barrier has a width of from 300  $\mu\text{m}$  to 30  $\mu\text{m}$ , and the distance between the center of each physical barrier to  
20 the center of adjacent physical barriers is at least 325  $\mu\text{m}$ .

In a preferred embodiment, a hydrophobic material is deposited onto the substrate to form barriers of the desired width using an ink-jet head, coupled to an appropriate robotic system. For example a microdrop dosing head, that has been adapted to apply a suspension or solution of a desired hydrophobic material (e.g., an oil based material that  
25 forms a barrier after the solvent has evaporated), may be coupled with an anorad gantry system and fitted to an appropriate housing and dispensing system so that a grid of the hydrophobic material may be applied onto the desired substrate forming a plurality of wells on the substrate. After the grid of hydrophobic material has been formed, different probes are spotted onto each well (or mixtures of probes may be applied to each well)  
30 using a robotic system similar to that used to form the grid, but that has been adapted to apply solutions or suspensions of probes. In one embodiment, the same robotic system is used to apply the hydrophobic grid and the probes. In this embodiment, the dispensing

WO 99/09217

PCT/US98/16966

system is flushed after the hydrophobic grid is applied and then primed for delivery of probe.

### **EXAMPLE 6**

#### **5 Preparation of Support Bound Oligonucleotides**

Oligonucleotides, i.e., small nucleic acid segments, may be readily prepared by, for example, directly synthesizing the oligonucleotide by chemical means, as is commonly practiced using an automated oligonucleotide synthesizer.

In general, oligonucleotides may be bound to a support through appropriate  
10 reactive groups. Such groups are well known in the art and include, for example, amino (-NH<sub>2</sub>); hydroxyl (-OH); or carboxyl (CO<sub>2</sub>H) groups. Support bound oligonucleotides may be prepared by any of the methods known to those of skill in the art using any suitable support such as glass, polystyrene or Teflon. One strategy is to precisely spot oligonucleotides synthesized by standard synthesizers. Immobilization can be achieved by  
15 many methods, including, for example, using passive adsorption (Inouye & Hondo, 1990); using UV light (Nagata *et al.*, 1985; Dahlen *et al.*, 1987; Morriey & Collins, 1989); or by covalent binding of base modified DNA (Keller *et al.*, 1988; 1989); or by formation of amide groups between the probe and the support (Wall *et al.*, 1995; Chebab *et al.*, 1992; and Zhang *et al.*, 1991); all references being specifically incorporated  
20 herein.

Another strategy that may be employed is the use of the strong biotin-streptavidin interaction as a linker. For example, Broude *et al.* (1994) describe the use of Biotinylated probes, although these are duplex probes, that are immobilized on streptavidin-coated magnetic beads. Streptavidin-coated beads may be purchased from  
25 Dynal, Oslo. Of course, this same linking chemistry is applicable to coating any surface with streptavidin. Biotinylated probes may be purchased from various sources, such as, e.g., Operon Technologies (Alameda, CA).

Nunc Laboratories (Naperville, IL) is also selling suitable material that could be used. Nunc Laboratories have developed a method by which DNA can be covalently  
30 bound to the microwell surface termed CovaLink NH. CovaLink NH is a polystyrene surface grafted with secondary amino groups (>NH) that serve as bridge-heads for further covalent coupling. CovaLink Modules may be purchased from Nunc

WO 99/09217

PCT/US98/16966

Laboratories. DNA molecules may be bound to CovaLink exclusively at the 5'-end by a phosphoramidate bond, allowing immobilization of more than 1 pmol of DNA (Rasmussen *et al.*, 1991).

The use of CovaLink NH strips for covalent binding of DNA molecules at the 5' 5'-end has been described (Rasmussen *et al.*, 1991). In this technology, a phosphoramidate bond is employed (Chu *et al.*, 1983). This is beneficial as immobilization using only a single covalent bond is preferred. The phosphoramidate bond joins the DNA to the CovaLink NH secondary amino groups that are positioned at the end of spacer arms covalently grafted onto the polystyrene surface through a 2 nm long spacer arm. To link an oligonucleotide to CovaLink NH via an phosphoramidate bond, the oligonucleotide terminus must have a 5'-end phosphate group. It is, perhaps, even possible for biotin to be covalently bound to CovaLink and then streptavidin used to bind the probes.

More specifically, the linkage method includes dissolving DNA in water (7.5 ng/ul) and denaturing for 10 min. at 95°C and cooling on ice for 10 min. Ice-cold 0.1 M 1-methylimidazole, pH 7.0 (1-MeIm<sub>7</sub>), is then added to a final concentration of 10 mM 1-MeIm<sub>7</sub>. A ss DNA solution is then dispensed into CovaLink NH strips (75 ul/well) standing on ice.

Carbodiimide 0.2 M 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (EDC), dissolved in 10 mM 1-MeIm<sub>7</sub>, is made fresh and 25 ul added per well. The strips are incubated for 5 hours at 50°C. After incubation the strips are washed using, e.g., Nunc-Immuno Wash; first the wells are washed 3 times, then they are soaked with washing solution for 5 min., and finally they are washed 3 times (where in the washing solution is 0.4 N NaOH, 0.25% SDS heated to 50°C).

It is contemplated that a further suitable method for use with the present invention is that described in PCT Patent Application WO 90/03382 (Southern & Maskos), incorporated herein by reference. This method of preparing an oligonucleotide bound to a support involves attaching a nucleoside 3'-reagent through the phosphate group by a covalent phosphodiester link to aliphatic hydroxyl groups carried by the support. The oligonucleotide is then synthesized on the supported nucleoside and protecting groups removed from the synthetic oligonucleotide chain under standard conditions that do not

WO 99/09217

PCT/US98/16966

cleave the oligonucleotide from the support. Suitable reagents include nucleoside phosphoramidite and nucleoside hydrogen phosphate.

An on-chip strategy for the preparation of DNA probe for the preparation of DNA probe arrays may be employed. For example, addressable laser-activated  
5 photodeprotection may be employed in the chemical synthesis of oligonucleotides directly on a glass surface, as described by Fodor *et al.* (1991), incorporated herein by reference. Probes may also be immobilized on nylon supports as described by Van Ness *et al.* (1991); or linked to Teflon using the method of Duncan & Cavalier (1988); all references being specifically incorporated herein.

10 To link an oligonucleotide to a nylon support, as described by Van Ness *et al.* (1991), requires activation of the nylon surface via alkylation and selective activation of the 5'-amine of oligonucleotides with cyanuric chloride.

One particular way to prepare support bound oligonucleotides is to utilize the light-generated synthesis described by Pease *et al.*, (1994, incorporated herein by  
15 reference). These authors used current photolithographic techniques to generate arrays of immobilized oligonucleotide probes (DNA chips). These methods, in which light is used to direct the synthesis of oligonucleotide probes in high-density, miniaturized arrays, utilize photolabile 5'-protected *N*-acyl-deoxynucleoside phosphoramidites, surface linker chemistry and versatile combinatorial synthesis strategies. A matrix of 256 spatially  
20 defined oligonucleotide probes may be generated in this manner and then used in the advantageous Format 3 sequencing, as described herein.

Of course, one could easily purchase a DNA chip, such as one of the light-activated chips described above, from a commercial source. In this regard, one may contact Affymetrix of Santa Clara, CA 95051, and Beckman.

25 In a preferred embodiment, the probes of the invention include an informational portion (the portion which hybridizes to the target nucleic acid and gives sequence information) a reactive group to be attached to the substrate (solid support), and randomized positions, i.e., any of the four bases may be found at these positions. A preferred probe has the sequence 5'-(T)<sub>6</sub>-(N)<sub>3</sub>-(B)<sub>5</sub>, where T = thymine (binds to solid  
30 support), N = A, C, G, or T (randomized positions), and B = the five information positions of the probe (informational portion). In a most preferred embodiment, the probe may be bound to the support and a spacer moiety is found at the end of the probe



WO 99/09217

PCT/US98/16966

or internal to the probe and 5' of (N)<sub>3</sub>. The spacers may be comprised of atoms capable of forming at least two covalent bonds such as carbon, silicon, oxygen, sulfur, phosphorous, and the like, or may be comprised of molecules capable of forming at least two covalent bonds such as sugar-phosphate groups, amino acids, peptides, nucleosides, nucleotides, sugars, carbohydrates, aromatic rings, hydrocarbon rings, linear and branched hydrocarbons, and the like.

### **EXAMPLE 7**

#### **Preparation of Nucleic Acid Fragments**

10 The nucleic acids to be sequenced may be obtained from any appropriate source, such as cDNAs, genomic DNA, chromosomal DNA, microdissected chromosome bands, cosmid or YAC inserts, and RNA, including mRNA without any amplification steps. For example, Sambrook *et al.* (1989) describes three protocols for the isolation of high molecular weight DNA from mammalian cells (p. 9.14-9.23).

15 Target nucleic acid fragments may be prepared as clones in M13, plasmid or lambda vectors and/or prepared directly from genomic DNA or cDNA by PCR or other amplification methods. Samples may be prepared or dispensed in multiwell plates. About 100-1000 ng of DNA samples may be prepared in 2-500 ml of final volume. Target nucleic acids prepared by PCR may be directly applied to a substrate for Format I SBH  
20 without purification. Once the target nucleic acids are fixed to the substrate, the substrate may be washed or directly annealed with probes.

The nucleic acids would then be fragmented by any of the methods known to those of skill in the art including, for example, using restriction enzymes as described at 9.24-9.28 of Sambrook *et al.* (1989), shearing by ultrasound and NaOH treatment.

25 Low pressure shearing is also appropriate, as described by Schrieffer *et al.* (1990, incorporated herein by reference). In this method, DNA samples are passed through a small French pressure cell at a variety of low to intermediate pressures. A lever device allows controlled application of low to intermediate pressures to the cell. The results of these studies indicate that low-pressure shearing is a useful alternative to sonic and  
30 enzymatic DNA fragmentation methods.

One particularly suitable way for fragmenting DNA is contemplated to be that using the two base recognition endonuclease, CviJI, described by Fitzgerald *et al.* (1992).

WO 99/09217

PCT/US98/16966

These authors described an approach for the rapid fragmentation and fractionation of DNA into particular sizes that they contemplated to be suitable for shotgun cloning and sequencing. The present inventor envisions that this will also be particularly useful for generating random, but relatively small, fragments of DNA for use in the present  
5 sequencing technology.

The restriction endonuclease *Cvi*JI normally cleaves the recognition sequence PuGCPy between the G and C to leave blunt ends. Atypical reaction conditions, which alter the specificity of this enzyme (*Cvi*JI\*\*), yield a quasi-random distribution of DNA fragments from the small molecule pUC19 (2688 base pairs). Fitzgerald *et al.* (1992)  
10 quantitatively evaluated the randomness of this fragmentation strategy, using a *Cvi*JI\*\* digest of pUC19 that was size fractionated by a rapid gel filtration method and directly ligated, without end repair, to a lac Z minus M13 cloning vector. Sequence analysis of 76 clones showed that *Cvi*JI\*\* restricts pyGCPy and PuGCPu, in addition to PuGCPy sites, and that new sequence data is accumulated at a rate consistent with random  
15 fragmentation.

As reported in the literature, advantages of this approach compared to sonication and agarose gel fractionation include: smaller amounts of DNA are required (0.2-0.5 ug instead of 2-5 ug); and fewer steps are involved (no preligation, end repair, chemical extraction, or agarose gel electrophoresis and elution are needed). These advantages are  
20 also proposed to be of use when preparing DNA for sequencing by Format 3.

In a preferred embodiment, the "fragments" of the nucleic acid sample are prepared so that they cannot be ligated to each other. Such a pool of fragments may be obtained by treating the fragmented nucleic acids obtained by enzyme digestion or physical shearing, with a phosphatase (e.g., calf intestinal phosphatase). Alternatively,  
25 nonligatable fragments of the sample nucleic acid may be obtained by using random primers (e.g., N<sub>5</sub>-N<sub>9</sub>, where N = A, G, T, or C), which have no phosphate at their 5'-ends, in a Sanger-dideoxy sequencing reaction with the sample nucleic acid. This will produce fragments of DNA that have a complementary sequence to the target nucleic acid and that are terminated in a dideoxy residue and which cannot be ligated to other  
30 fragments.

Irrespective of the manner in which the nucleic acid fragments are obtained or prepared, it is important to denature the DNA to give single stranded pieces available for

WO 99/09217

PCT/US98/16966

hybridization. This is achieved by incubating the DNA solution for 2-5 minutes at 80-90°C. The solution is then cooled quickly to 2°C to prevent renaturation of the DNA fragments before they are contacted with the chip.

## 5 **EXAMPLE 8**

### **Preparation of DNA Arrays**

Arrays may be prepared by spotting DNA samples on a support such as a nylon membrane. Spotting may be performed by using arrays of metal pins (the positions of which correspond to an array of wells in a microtiter plate) to repeated by transfer of  
10 about 20 nl of a DNA solution to a nylon membrane. By offset printing, a density of dots higher than the density of the wells is achieved. One to 25 dots may be accommodated in 1 mm<sup>2</sup>, depending on the type of label used. By avoiding spotting in some preselected number of rows and columns, separate subsets (subarrays) may be formed. Samples in one subarray may be the same genomic segment of DNA (or the same gene) from  
15 different individuals, or may be different, overlapped genomic clones. Each of the subarrays may represent replica spotting of the same samples. In one example, a selected gene segment may be amplified from 64 patients. For each patient, the amplified gene segment may be in one 96-well plate (all 96 wells containing the same sample). A plate for each of the 64 patients is prepared. By using a 96-pin device, all samples may be  
20 spotted on one 8 x 12 cm membrane. Subarrays may contain 64 samples, one from each patient. Where the 96 subarrays are identical, the dot span may be 1 mm<sup>2</sup> and there may be a 1 mm space between subarrays.

Another approach is to use membranes or plates (available from NUNC, Naperville, Illinois) which may be partitioned by physical spacers e.g. a plastic grid  
25 molded over the membrane, the grid being similar to the sort of membrane applied to the bottom of multiwell plates, or hydrophobic strips. A fixed physical spacer is not preferred for imaging by exposure to flat phosphor-storage screens or x-ray films.

## **EXAMPLE 9**

### 30 **Hybridization and Scoring Process**

Labeled probes may be mixed with hybridization buffer and pipetted, preferably by multichannel pipettes, to the subarrays. To prevent mixing of the probes between

WO 99/09217

PCT/US98/16966

subarrays (if there are no hydrophobic strips or physical barriers imprinted in the membrane), a corresponding plastic, metal or ceramic grid may be firmly pressed to the membrane. Also, the volume of the buffer may be reduced to about 1 ml or less per mm<sup>2</sup>. The concentration of the probes and hybridization conditions used may be as described

5 previously except that the washing buffer may be quickly poured over the array of subarrays to allow fast dilution of probes and thus prevent significant cross-hybridization. For the same reason, a minimal concentration of the probes may be used and hybridization time extended to the maximal practical level. For DNA detection and sequencing, knowledge of a "normal" sequence allows the use of the continuous stacking

10 interaction phenomenon to increase the signal. In addition to the labelled probe, additional unlabelled probes which hybridize back to back with a labelled one may be added in the hybridization reaction. The amount of the hybrid may be increased several times. The probes may be connected by ligation. This approach may be important for resolving DNA regions forming "compressions".

15 In the case of radiolabelled probes, images of the filters may be obtained, preferably by phosphorstorage technology. Fluorescent labels may be scored by CCD cameras, confocal microscopy or otherwise. In order to properly scale and integrate data from different hybridization experiments, raw signals are normalized based on the amount of target in each dot. Differences in the amount of target DNA per dot may be corrected

20 for by dividing signals of each probe by an average signal for all probes scored on one dot. The normalized signals may be scaled, usually from 1-100, to compare data from different experiments. Also, in each subarray, several control DNAs may be used to determine an average background signal in those samples which do not contain a full match target. For samples obtained from diploid (polyploid) scores, homozygotic controls

25 may be used to allow recognition of heterozygotes in the samples.

### **EXAMPLE 10**

#### **Hybridization With Oligonucleotides**

Oligonucleotides were either purchased from Genosys Inc., Houston, Texas or

30 made on an Applied Biosystems 381A DNA synthesizer. Most of the probes used were not purified by HPLC or gel electrophoresis. For example, probes were designed to have both a single perfectly complementary target in interferon, a M13 clone containing a 921

WO 99/09217

PCT/US98/16966

bp Eco RI-Bgl II human B1 - interferon fragment (Ohno and Tangiuchi, Proc. Natl. Acad. Sci. 74: 4370-4374 (1981)), and at least one target with an end base mismatch in M13 vector itself.

End labeling of oligonucleotides was performed as described [Maniatis et al.,

- 5 *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Cold Spring Harbor, New York (1982)] in 10 ml containing T4-polynucleotide kinase (5 units Amersham),  $\gamma^{32}\text{P}$ -ATP (3.3 pM, 10 mCi Amersham 3000 Ci/mM) and oligonucleotide (4 pM, 10 ng). Specific activities of the probes were  $2.5\text{-}5 \times 10^9$  cpm/nM.

- Single stranded DNA (2 to 4 ml in 0.5 NaOH, 1.5 M NaCl) was spotted on a  
10 Gene Screen membrane wetted with the same solution, the filters were neutralized in 0.05 M  $\text{Na}_2\text{HPO}_4$  pH 6.5, baked in an oven at 80°C for 60 min. and UV irradiated for 1 min. Then, the filters were incubated in hybridization solution (0.5 M  $\text{Na}_2\text{HPO}_4$  pH 7.2, 7% sodium lauroyl sarcosine for 5 min at room temperature and placed on the surface of a plastic Petri dish. A drop of hybridization solution (10 ml, 0.5 M  $\text{Na}_2\text{HPO}_4$  pH 7.2, 7%  
15 sodium lauroyl sarcosine) with a  $^{32}\text{P}$  end-labeled oligomer probe at 4 nM concentration was placed over 1-6 dots per filter, overlaid with a square piece of polyethylene (approximately 1 X 1 cm.), and incubated in a moist chamber at the indicated temperatures for 3 hr. Hybridization was stopped by placing the filter in 6X SSC washing solution for 3 X 5 minute at 0°C to remove unhybridized probe. The filter was  
20 either dried, or further washed for the indicated times and temperatures, and autoradiographed. For discrimination measurements, the dots were excised from the dried filters after autoradiography [a phosphorimager (Molecular Dynamics, Sunnyvale, California) may be used] placed in liquid scintillation cocktail and counted. The uncorrected ratio of cpm for IF and M13 dots is given as D.

- 25 The conditions reported herein allow hybridization with very short oligonucleotides but ensure discriminations between matched and mismatched oligonucleotides that are complementary to and therefore bind to a target nucleic acid. Factors which influence the efficient detection of hybridization of specific short sequences based on the degree of discriminations (D) between a perfectly complementary target and an imperfectly  
30 complementary target with a single mismatch in the hybrid are defined. In experimental tests, dot blot hybridization of twenty-eight probes that were 6 to 8 nucleotides in length

WO 99/09217

PCT/US98/16966

to two M13 clones or to model oligonucleotides bound to membrane filters was accomplished. The principles guiding the experimental procedures are given below.

Oligonucleotide hybridization to filter bound target nucleic acids only a few nucleotides longer than the probe in conditions of probe excess is a pseudo-first order reaction with respect to target concentration. This reaction is defined by:

$$S_t/S_o = e^{-k_h [OP] t}$$

Wherein  $S_t$  and  $S_o$  are target sequence concentrations at time  $t$  and  $t_o$ , respectively.  $(OP)$  is probe concentration and  $t$  is temperature. The rate constant for hybrid formation,  $k_h$ , increases only slightly in the 0°C to 30°C range (Porschke and Eigen, *J. Mol. Biol.* 62: 361 (1971); Craig et al., *J. Mol. Biol.* 62: 383 (1971)). Hybrid melting is a first order reaction with respect to hybrid concentration (here replaced by mass due to filter bound state) as shown in:

$$H_t/H_o = e^{-k_m t}$$

In this equation,  $H_t$  and  $H_o$  are hybrid concentrations at times  $t$  and  $t_o$ , respectively;  $k_m$  is a rate constant for hybrid melting which is dependent on temperature and salt concentration [Ikuta et al., *Nucl. Acids Res.* 15: 797 (1987); Porschke and Eigen, *J. Mol. Biol.* 62: 361 (1971); Craig et al., *J. Mol. Biol.* 62: 303 (1971)]. During hybridization, which is a strand association process, the back, melting, or strand dissociation, reaction takes place as well. Thus, the amount of hybrid formed in time is result of forward and back reactions. The equilibrium may be moved towards hybrid formation by increasing probe concentration and/or decreasing temperature. However, during washing cycles in large volumes of buffer, the melting reaction is dominant and the back reaction hybridization is insignificant, since the probe is absent. This analysis indicates workable Short Oligonucleotide Hybridization (SOH) conditions can be varied for probe concentration or temperature.

D or discrimination is defined in equation four:

$$D = H_p(t_w) / H_i(t_w)$$

$H_p(t_w)$  and  $H_i(t_w)$  are the amounts hybrids remaining after a washing time,  $t_w$ , for the identical amounts of perfectly and imperfectly complementary duplex, respectively.

For a given temperature, the discrimination  $D$  changes with the length of washing time and reaches the maximal value when  $H_i = B$  which is equation five.

WO 99/09217

PCT/US98/16966

The background, B, represents the lowest hybridization signal detectable in the system. Since any further decrease of  $H_i$  may not be examined, D increases upon continued washing. Washing past  $t_w$  just decreases  $H_p$  relative to B, and is seen as a decrease in D. The optimal washing time,  $t_w$ , for imperfect hybrids, from equation three  
 5 and equation five is:

$$t_w = -\ln(B / H_i(t_0)) / k_{m,i}$$

Since  $H_p$  is being washed for the same  $t_w$ , combining equations, one obtains the optimal discrimination function:

$$D = e^{\ln(B/H_i(t_0)) k_{m,p} / k_{m,i}} \times H_p(t_0) / B$$

10 The change of D as a function, of T is important because of the choice of an optimal washing temperature. It is obtained by substituting the Arrhenius equation which is:

$$K = A e^{-E_a / RT}$$

into the previous equation to form the final equation:

$$15 \quad D = H_p(t_0) / B \times (B / H_i(t_0))^{(A_p / A_i) e^{(E_{a,i} - E_{a,p}) / RT}};$$

Wherein B is less than  $H_i(t_0)$ .

Since the activation energy for perfect hybrids,  $E_{a,p}$ , and the activation energy for imperfect hybrids,  $E_{a,i}$ , can be either equal, or  $E_{a,i}$  less than  $E_{a,p}$ , D is temperature independent, or decreases with increasing temperature, respectively. This result implies  
 20 that the search for stringent temperature conditions for good discrimination in SOH is unjustified. By washing at lower temperatures, one obtains equal or better discrimination, but the time of washing exponentially increases with the decrease of temperature. Discrimination more strongly decreases with T, if  $H_i(t_0)$  increases relative to  $H_p(t_0)$ .

D at lower temperatures depends to a higher degree on the  $H_p(t_0) / B$  ratio than on  
 25 the  $H_p(t_0) / H_i(t_0)$  ratio. This result indicates that it is better to obtain a sufficient quantity of  $H_p$  in the hybridization regardless of the discrimination that can be achieved in this step. Better discrimination can then be obtained by washing, since the higher amounts of perfect hybrid allow more time for differential melting to show an effect. Similarly, using larger amounts of target nucleic acid a necessary discrimination can be  
 30 obtained even with small differences between  $K_{m,p}$  and  $K_{m,i}$ .

Extrapolated to a more complex situation than covered in this simple model, the result is that washing at lower temperatures is even more important for obtaining

WO 99/09217

PCT/US98/16966

discrimination in the case of hybridization of a probe having many end-mismatches within a given nucleic acid target.

Using the described theoretical principles as a guide for experiments, reliable hybridizations have been obtained with probes six to eight nucleotides in length. All experiments were performed with a floating plastic sheet providing a film of hybridization solution above the filter. This procedure allows maximal reduction in the amount of probe, and thus reduced label costs in dot blot hybridizations. The high concentration of sodium lauroyl sarcosine instead of sodium lauroyl sulfate in the phosphate hybridization buffer allows dropping the reaction from room temperature down to 12°C. Similarly, the 4-6 X SSC, 10% sodium lauroyl sarcosine buffer allows hybridization at temperatures as low as 2°C. The detergent in these buffers is for obtaining tolerable background with up to 40 nM concentrations of labelled probe. Preliminary characterization of the thermal stability of short oligonucleotide hybrids was determined on a prototype octamer with 50% G+C content, i.e. probe of sequence TGCTCATG. The theoretical expectation is that this probe is among the less stable octamers. Its transition enthalpy is similar to those of more stable heptamers or, even to probes 6 nucleotides in length (Bresslauer et al., *Proc. Natl. Acad. Sci. U.S.A.* **83**: 3746 (1986)). Parameter  $T_d$ , the temperature at which 50% of the hybrid is melted in unit time of a minute is 18°C. The result shows that  $T_d$  is 15°C lower for the 8 bp hybrid than for an 11 bp duplex [Wallace et al., *Nucleic Acids Res.* **6**: 3543 (1979)].

In addition to experiments with model oligonucleotides, an M13 vector was chosen as a system for a practical demonstration of short oligonucleotide hybridization. The main aim was to show useful end-mismatch discrimination with a target similar to the ones which will be used in various applications of the method of the invention. Oligonucleotide probes for the M13 model were chosen in such a way that the M13 vector itself contains the end mismatched base. Vector IF, an M13 recombinant containing a 921 bp human interferon gene insert, carries single perfectly matched target. Thus, IF has either the identical or a higher number of mismatched targets in comparison to the M13 vector itself.

Using low temperature conditions and dot blots, sufficient differences in hybridization signals were obtained between the dot containing the perfect and the mismatched targets and the dot containing the mismatched targets only. This was true for



WO 99/09217

PCT/US98/16966

the 6-mer oligonucleotides and was also true for the 7 and 8-mer oligonucleotides hybridized to the large IF-M13 pair of nucleic acids.

The hybridization signal depends on the amount of target available on the filter for reaction with the probe. A necessary control is to show that the difference in sign  
5 intensity is not a reflection of varying amounts of nucleic acid in the two dots. Hybridization with a probe that has the same number and kind of targets in both IF and M13 shows that there is an equal amount of DNA in the dots. Since the efficiency of hybrid formation increases with hybrid length, the signal for a duplex having six nucleotides was best detected with a high mass of oligonucleotide target bound to the  
10 filter. Due to their lower molecular weight, a larger number of oligonucleotide target molecules can be bound to a given surface area when compared to large molecules of nucleic acid that serves as target.

To measure the sensitivity of detection with unpurified DNA, various amounts of phage supernatants were spotted on the filter and hybridized with a  $^{32}$  P-labelled octamer.  
15 As little as 50 million unpurified phage containing no more than 0.5 ng of DNA gave a detectable signal indicating that sensitivity of the short oligonucleotide hybridization method is sufficient. Reaction time is short, adding to the practicality.

As mentioned in the theoretical section above, the equilibrium yield of hybrid depends on probe concentration and/or temperature of reaction. For instance, the signal  
20 level for the same amount of target with 4 nM octamer at 13°C is 3 times lower than with a probe concentration of 40 nM, and is decreased 4.5-times by raising the hybridization temperature to 25°C.

The utility of the low temperature wash for achieving maximal discrimination is demonstrated. To make the phenomenon visually obvious, 50 times more DNA was put in  
25 the M13 dot than in the IF dot using hybridization with a vector specific probe. In this way, the signal after the hybridization step with the actual probe was made stronger in the, mismatched than in the matched case. The  $H_p/H_i$  ratio was 1:4. Inversion of signal intensities after prolonged washing at 7°C was achieved without a massive loss of perfect hybrid, resulting in a ratio of 2:1. In contrast, it is impossible to achieve any  
30 discrimination at 25°C, since the matched target signal is already brought down to the background level with 2 minute washing; at the same time, the signal from the mismatched hybrid is still detectable. The loss of discrimination at 13°C compared to 7°C

WO 99/09217

PCT/US98/16966

is not so great but is clearly visible. If one considers the 90 minute point at 7°C and the 15 minute point at 13°C when, the mismatched hybrid signal is near the background level, which represents optimal washing times for the respective conditions, it is obvious that the amount of several times greater at 7°C than at 13°C. To illustrate this further, 5 the time course of the change discrimination with washing of the same amount of starting hybrid at the two temperatures shows the higher maximal D at the lower temperature. These results confirm the trend in the change of D with temperature and the ratio of amounts of the two types of hybrid at the start of the washing step.

In order to show the general utility of the short oligonucleotide hybridization 10 conditions, we have looked hybridization of 4 heptamers, 10 octamers and an additional 14 probes up to 12 nucleotides in length in our simple M13 system. These include the nonamer GTTTTTTAA and octamer GGCAGGCG representing the two extremes of GC content. Although GC content and sequence are expected to influence the stability of short hybrids [Bresslauer et al., *Proc. Natl. Acad. Sci. U.S.A.* 83: 3746 (1986)], the 15 low temperature short oligonucleotide conditions were applicable to all tested probes in achieving sufficient discrimination. Since the best discrimination value obtained with probes 13 nucleotides in length was 20, a several fold drop due to sequence variation is easily tolerated.

The M13 system has the advantage of showing the effects of target DNA 20 complexity on the levels of discrimination. For two octamers having either none or five mismatched targets and differing in only one GC pair the observed discriminations were 18.3 and 1.7, respectively.

In order to show the utility of this method, three probes 8 nucleotides in length were tested on a collection of 51 plasmid DNA dots made from a library in Bluescript 25 vector. One probe was present and specific for Bluescript vector but was absent in M13, while the other two probes had targets that were inserts of known sequence. This system allowed the use of hybridization negative or positive control DNAs with each probe. This probe sequence (CTCCCTTT) also had a complementary target in the interferon insert. Since the M13 dot is negative while the interferon insert in either M13 or Bluescript was 30 positive, the hybridization is sequence specific. Similarly, probes that detect the target sequence in only one of 51 inserts, or in none of the examined inserts along with controls

WO 99/09217

PCT/US98/16966

that confirm that hybridization would have occurred if the appropriate targets were present in the clones.

- Thermal stability curves for very short oligonucleotide hybrids that are 6-8 nucleotides in length are at least 15°C lower than for hybrids 11-12 nucleotides in length [Fig. 1 and Wallace et al., *Nucleic Acids Res.* 6: 3543-3557 (1979)]. However, performing the hybridization reaction at a low temperature and with a very practical 0.4-40 nM concentration of oligonucleotide probe allows the detection of complementary sequence in a known or unknown nucleic acid target. To determine an unknown nucleic acid sequence completely, an entire set containing 65,535 8-mer probes may be used.
- 10 Sufficient amounts of nucleic acid for this purpose are present in convenient biological samples such as a few microliters of M13 culture, a plasmid prep from 10 ml of bacterial culture or a single colony of bacteria, or less than 1 ml of a standard PCR reaction.

- Short oligonucleotides 6-10 nucleotides long give excellent discrimination. The relative decrease in hybrid stability with a single end mismatch is greater than for longer probes. Results with the octamer TGCTCATG support this conclusion. In the experiments, the target with a G/T end mismatch, hybridization to the target of this type of mismatch is the most stable of all other types of oligonucleotide. This discrimination achieved is the same as or greater than an internal G/T mismatch in a 19 base paired duplex greater than an internal G/T mismatch in a 19 paired duplex [Ikuta et al., *Nucl. Acids res.* 15: 797 (1987)]. Exploiting these discrimination properties using the described hybridization conditions for short oligonucleotide hybridization allows a very precise determination of oligonucleotide targets. In contrast to the ease of detecting discrimination between perfect and imperfect hybrids, a problem that may exist with using very short oligonucleotides is the preparation of sufficient amounts of hybrids. In practice, the need to discriminate  $H_p$  and  $H_i$  is aided by increasing the amount of DNA in the dot and/or the probe concentration, or by decreasing the hybridization temperature. However, higher probe concentrations usually increase background. Moreover, there are limits to the amounts of target nucleic acid that are practical to use. This problems was solved by the higher concentration of the detergent Sarcosyl which gave an effective background with 4 nM of probe. Further improvements may be effected either in the use of competitors for unspecific binding of probe to filter, or by changing the hybridization support material. Moreover, for probes having  $E_a$  less than 45 Kcal/mol (e.g. for many heptamers and a
- 25  
30

WO 99/09217

PCT/US98/16966

majority of hexamers, modified oligonucleotides give a more stable hybrid [Asseline, et al., *Proc. Nat'l Acad. Sci.* 81: 3297 (1984)] than their unmodified counterparts. The hybridization conditions described in this invention for short oligonucleotide hybridization using low temperatures give better discriminating for all sequences and duplex hybrid  
5 inputs. The only price paid in achieving uniformity in hybridization conditions for different sequences is an increase in washing time from minutes to up to 24 hours depending on the sequence. Moreover, the washing time can be further reduced by decreasing the salt concentration.

Although there is excellent discrimination of one matched hybrid over a  
10 mismatched hybrids, in short oligonucleotide hybridization, signals from mismatched hybrids exist, with the majority of the mismatch hybrids resulting from end mismatch. This may limit insert sizes that may be effectively examined by a probe of a certain length.

The influence of sequence complexity on discrimination cannot be ignored.  
15 However, the complexity effects are more significant when defining sequence information by short oligonucleotide hybridization for specific, nonrandom sequences, and can be overcome by using an appropriate probe to target length ratio. The length ratio is chosen to make unlikely, on statistical grounds, the occurrence of specific sequences which have a number of end-mismatches which would be able to eliminate or falsely invert  
20 discrimination. Results suggest the use of oligonucleotides 6, 7, and 8 nucleotides in length on target nucleic acid inserts shorter than 0.6, 2.5, and 10 kb, respectively.

### EXAMPLE 11

#### DNA Sequencing

25 An array of subarrays allows for efficient sequencing of a small set of samples arrayed in the form of replicated subarrays; For example, 64 samples may be arrayed on a 8 X 8 mm subarray and 16 X 24 subarrays may be replicated on a 15 X 23 cm membrane with 1 mm wide spacers between the subarrays. Several replica membranes may be made. For example, probes from a universal set of three thousand seventy-two  
30 7-mers may be divided in thirty-two 96-well plates and labelled by kinasing. Four membranes may be processed in parallel during one hybridization cycle. On each membrane, 384 probes may be scored. All probes may be scored in two hybridization

WO 99/09217

PCT/US98/16966

cycles. Hybridization intensities may be scored and the sequence assembled as described below.

If a single sample subarray or subarrays contains several unknowns, especially when similar samples are used, a smaller number of probes may be sufficient if they are  
5 intelligently selected on the basis of results of previously scored probes. For example, if probe AAAAAAA is not positive, there is a small chance that any of 8 overlapping probes are positive. If AAAAAAA is positive, then two probes are usually positive. The sequencing process in this case consists of first hybridizing a subset of minimally  
10 overlapped probes to define positive anchors and then to successively select probes which confirms one of the most likely hypotheses about the order of anchors and size and type of gaps between them. In this second phase, pools of 2-10 probes may be used where each probe is selected to be positive in only one DNA sample which is different from the samples expected to be positive with other probes from the pool.

The subarray approach allows efficient implementation of probe competition  
15 (overlapped probes) or probe cooperation (continuous stacking of probes) in solving branching problems. After hybridization of a universal set of probes the sequence assembly program determines candidate sequence subfragments (Sfs). For the further assembly of Sfs, additional information has to be provided (from overlapped sequences of DNA fragments, similar sequences, single pass gel sequences, or from other hybridization  
20 or restriction mapping data). Primers for single pass gel sequencing through the branch points are identified from the SBH sequence information or from known vector sequences, e.g., the flanking sequences to the vector insert site, and standard Sanger-sequencing reactions are performed on the sample DNA. The sequence obtained from this single pass gel sequencing is compared to the Sfs that read into and out of the branch  
25 points to identify the order of the Sfs. Further, single pass gel sequencing may be combined with SBH to de novo sequence or re-sequence a nucleic acid.

Competitive hybridization and continuous stacking interactions can also be used to assemble Sfs. These approaches are of limited commercial value for sequencing of large numbers of samples by SBH wherein a labelled probe is applied to a sample affixed to an  
30 array if a uniform array is used. Fortunately, analysis of small numbers of samples using replica subarrays allows efficient implementation of both approaches. On each of the replica subarrays, one branching point may be tested for one or more DNA samples using

WO 99/09217

PCT/US98/16966

pools of probes similarly as in solving mutated sequences in different samples spotted in the same subarray (see above).

If in each of 64 samples described in this example, there are about 100 branching points, and if 8 samples are analyzed in parallel in each subarray, then at least 800 subarray probings solve all branches. This means that for the 3072 basic probings an additional 800 probings (25%) are employed. More preferably, two probings are used for one branching point. If the subarrays are smaller, less additional probings are used. For example, if subarrays consist of 16 samples, 200 additional probings may be scored (6%). By using 7-mer probes ( $N_{1-2}B_7N_{1-2}$ ) and competitive or collaborative branching solving approaches or both, fragments of about 1000 bp fragments may be assembled by about 4000 probings. Furthermore, using 8-mer probes ( $NB_8N$ ) 4 kb or longer fragments may be assembled with 12,000 probings. Gapped probes, for example,  $NB_4NB_3N$  or  $NB_4NB_4N$  may be used to reduce the number of branching points.

## 15 EXAMPLE 12

### DNA Analysis by Transient Attachment to Subarrays of Probes and Ligation of Labelled Probes

Oligonucleotide probes having an informative length of four to 40 bases are synthesized by standard chemistry and stored in tubes or in multiwell plates. Specific sets of probes comprising one to 10,000 probes are arrayed by deposition or in situ synthesis on separate supports or distinct sections of a larger support. In the last case, sections or subarrays may be separated by physical or hydrophobic barriers. The probe arrays may be prepared by in situ synthesis. A sample DNA of appropriate size is hybridized with one or more specific arrays. Many samples may be interrogated as pools at the same subarrays or independently with different subarrays within one support. Simultaneously with the sample or subsequently, a single labelled probe or a pool of labelled probes is added on each of the subarrays. If attached and labelled probes hybridize back to back on the complementary target in the sample DNA they are ligated. Occurrence of ligation will be measured by detecting a label from the probe.

30 This procedure is a variant of the described DNA analysis process in which DNA samples are not permanently attached to the support. Transient attachment is provided by probes fixed to the support. In this case there is no need for a target DNA arraying

WO 99/09217

PCT/US98/16966

process. In addition, ligation allows detection of longer oligonucleotide sequences by combining short labelled probes with short fixed probes.

The process has several unique features. Basically, the transient attachment of the target allows its reuse. After ligation occur the target may be released and the label will  
5 stay covalently attached to the support. This feature allows cycling the target and production of detectable signal with a small quantity of the target. Under optimal conditions, targets do not need to be amplified, e.g. natural sources of the DNA samples may be directly used for diagnostics and sequencing purposes. Targets may be released by cycling the temperature between efficient hybridization and efficient melting of duplexes.  
10 More preferably, there is no cycling. The temperature and concentrations of components may be defined to have an equilibrium between free targets and targets entered in hybrids at about 50:50% level. In this case there is a continuous production of ligated products. For different purposes different equilibrium ratios are optimal.

An electric field may be used to enhance target use. At the beginning, a horizontal  
15 field pulsing within each subarray may be employed to provide for faster target sorting. In this phase, the equilibrium is moved toward hybrid formation, and unlabelled probes may be used. After a target sorting phase, an appropriate washing (which may be helped by a vertical electric field for restricting movement of the samples) may be performed. Several cycles of discriminative hybrid melting, target harvesting by hybridization and  
20 ligation and removing of unused targets may be introduced to increase specificity. In the next step, labelled probes are added and vertical electrical pulses may be applied. By increasing temperature, an optimal free and hybridized target ratio may be achieved. The vertical electric field prevents diffusion of the sorted targets.

The subarrays of fixed probes and sets of labelled probes (specially designed or  
25 selected from a universal probe set) may be arranged in various ways to allow an efficient and flexible sequencing and diagnostics process. For example, if a short fragment (about 100-500 bp) of a bacterial genome is to be partially or completely sequenced, small arrays of probes (5-30 bases in length) designed on the bases of known sequence may be used. If interrogated with a different pool of 10 labelled probes per subarray, an array of 10  
30 subarrays each having 10 probes, allows checking of 200 bases, assuming that only two bases connected by ligation are scored. Under the conditions where mismatches are discriminated throughout the hybrid, probes may be displaced by more than one base to

WO 99/09217

PCT/US98/16966

cover the longer target with the same number of probes. By using long probes, the target may be interrogated directly without amplification or isolation from the rest of DNA in the sample. Also, several targets may be analyzed (screened for) in one sample simultaneously. If the obtained results indicate occurrence of a mutation (or a pathogen),

5 additional pools of probes may be used to detect type of the mutation or subtype of pathogen. This is a desirable feature of the process which may be very cost effective in preventive diagnosis where only a small fraction of patients is expected to have an infection or mutation.

In the processes described in the examples, various detection methods may be

10 used, for example, radiolabels, fluorescent labels, enzymes or antibodies (chemiluminescence), large molecules or particles detectable by light scattering or interferometric procedures.

### **EXAMPLE 13**

#### **15 Sequencing a Target Using Octamers and Nonamers**

Data resulting from the hybridization of octamer and nonamer oligonucleotides shows that sequencing by hybridization provides an extremely high degree of accuracy. In this experiment, a known sequence was used to predict a series of contiguous overlapping component octamer and nonamer oligonucleotides.

20 In addition to the perfectly matching oligonucleotides, mismatch oligonucleotides, mismatch oligonucleotides wherein internal or end mismatches occur in the duplex formed by the oligonucleotide and the target were examined. In these analyses, the lowest practical temperature was used to maximize hybridization formation. Washes were accomplished at the same or lower temperatures to ensure maximal discrimination by

25 utilizing the greater dissociation rate of mismatch versus matched oligonucleotide/target hybridization. These conditions are shown to be applicable to all sequences although the absolute hybridization yield is shown to be sequence dependent.

The least destabilizing mismatch that can be postulated is a simple end mismatch, so that the test of sequencing by hybridization is the ability to discriminate perfectly

30 matched oligonucleotide/target duplexes from end-mismatched oligonucleotide/target duplexes.



WO 99/09217

PCT/US98/16966

The discriminative values for 102 of 105 hybridizing oligonucleotides in a dot blot format were greater than 2 allowing a highly accurate generation of the sequence. This system also allowed an analysis of the effect of sequence on hybridization formation and hybridization instability.

- 5        One hundred base pairs of a known portion of a human-interferon genes prepared by PCR, i.e. a 100 bp target sequence, was generated with data resulting from the hybridization of 105 oligonucleotides probes of known sequence to the target nucleic acid. The oligonucleotide probes used included 72 octamer and 21 nonamer oligonucleotides whose sequence was perfectly complementary to the target. The set of 93 probes  
10       provided consecutive overlapping frames of the target sequence e displaced by one or two bases.

- To evaluate the effect of mismatches, hybridization was examined for 12 additional probes that contained at least one end mismatch when hybridized to the 100 bp test target sequence. Also tested was the hybridization of twelve probes with target end-mismatched  
15       to four other control nucleic acid sequences chosen so that the 12 oligonucleotides formed perfectly matched duplex hybrids with the four control DNAs. Thus, the hybridization of internal mismatched, end-mismatched and perfectly matched duplex pairs of oligonucleotide and target were evaluated for each oligonucleotide used in the experiment. The effect of absolute DNA target concentration on the hybridization with the test  
20       octamer and nonamer oligonucleotides was determined by defining target DNA concentration by detecting hybridization of a different oligonucleotide probe to a single occurrence non- target site within the co-amplified plasmid DNA.

- The results of this experiment showed that all oligonucleotides containing perfect matching complementary sequence to the target or control DNA hybridized more strongly  
25       than those oligonucleotides having mismatches. To come to this conclusion, we examined  $H_p$  and D values for each probe.  $H_p$  defines the amount of hybrid duplex formed between a test target and an oligonucleotide probe. By assigning values of between 0 and 10 to the hybridization obtained for the 105 probes, it was apparent that 68.5% of the 105 probes had an  $H_p$  greater than 2.

- 30       Discrimination (D) values were obtained where D was defined as the ratio of signal intensities between 1) the dot containing a perfect matched duplex formed between test oligonucleotide and target or control nucleic acid and 2) the dot containing a

WO 99/09217

PCT/US98/16966

mismatch duplex formed between the same oligonucleotide and a different site within the target or control nucleic acid. Variations in the value of D result from either 1) perturbations in the hybridization efficiency which allows visualization of signal over background, or 2) the type of mismatch found between the test oligonucleotide and the target. The D values obtained in this experiment were between 2 and 40 for 102 of the 105 oligonucleotide probes examined. Calculations of D for the group of 102 oligonucleotides as a whole showed the average D was 10.6.

There were 20 cases where oligonucleotide/target duplexes exhibited an end-mismatch. In five of these, D was greater than 10. The large D value in these cases is most likely due to hybridization destabilization caused by other than the most stable (G/T and G/A) end mismatches. The other possibility is there was an error in the sequence of either the oligonucleotides or the target.

Error in the target for probes with low  $H_p$  was excluded as a possibility because such an error would have affected the hybridization of each of the other eight overlapping oligonucleotides. There was no apparent instability due to sequence mismatch for the other overlapping oligonucleotides, indicating the target sequence was correct. Error in the oligonucleotide sequence was excluded as a possibility after the hybridization of seven newly synthesized oligonucleotides was re-examined. Only 1 of the seven oligonucleotides resulted in a better D value. Low hybrid formation values may result from hybrid instability or from an inability to form hybrid duplex. An inability to form hybrid duplexes would result from either 1) self complementarity of the chosen probe or 2) target/target self hybridization. Oligonucleotide/oligonucleotide duplex formation may be favored over oligonucleotide/target hybrid duplex formation if the probe was self-complementary. Similarly, target/target association may be favored if the target was self-complementary or may form internal palindromes. In evaluating these possibilities, it was apparent from probe analysis that the questionable probes did not form hybrids with themselves. Moreover, in examining the contribution of target/target hybridization, it was determined that one of the questionable oligonucleotide probes hybridized inefficiently with two different DNAs containing the same target. The low probability that two different DNAs have a self-complementary region for the same target sequence leads to the conclusion that target/target hybridization did not contribute to low hybridization formation. Thus, these results indicate that hybrid instability and not the

WO 99/09217

PCT/US98/16966

inability to form hybrids was the cause of the low hybrid formation observed for specific oligonucleotides. The results also indicate that low hybrid formation is due to the specific sequences of certain oligonucleotides. Moreover, the results indicate that reliable results may be obtained to generate sequences if octamer and nonamer oligonucleotides are used.

5        These results show that using the methods described long sequences of any specific target nucleic acid may be generated by maximal and unique overlap of constituent oligonucleotides. Such sequencing methods are dependent on the content of the individual component oligomers regardless of their frequency and their position.

      The sequence which is generated using the algorithm described below is of high  
10 fidelity. The algorithm tolerates false positive signals from the hybridization dots as is indicated from the fact the sequence generated from the 105 hybridization values, which included four less reliable values, was correct. This fidelity in sequencing by hybridization is due to the "all or none" kinetics of short oligonucleotide hybridization and the difference in duplex stability that exists between perfectly matched duplexes and  
15 mismatched duplexes. The ratio of duplex stability of matched and end-mismatched duplexes increases with decreasing duplex length. Moreover, binding energy decreases with decreasing duplex length resulting in a lower hybridization efficiency. However, the results provided show that octamer hybridization allows the balancing of the factors affecting duplex stability and discrimination to produce a highly accurate method of  
20 sequencing by hybridization. Results presented in other examples show that oligonucleotides that are 6, 7, or 8 nucleotides can be effectively used to generate reliable sequence on targets that are 0.5 kb (for hexamers) 2 kb (for septamers) and 6kb (for octamers). The sequence of long fragments may be overlapped to generate a complete genome sequence.

25

#### **EXAMPLE 14**

##### **Analyzing the Data Obtained**

Image files are analyzed by an image analysis program, like DOTS program (Drmanac *et al.*, 1993), and scaled and evaluated by statistical functions included, e.g., in  
30 SCORES program (Drmanac *et al.* 1994). From the distribution of the signals an optimal threshold is determined for transforming signal into +/- output. From the position of the label detected, F + P nucleotide sequences from the fragments would be determined by

WO 99/09217

PCT/US98/16966

combining the known sequences of the immobilized and labeled probes corresponding to the labeled positions. The complete nucleic acid sequence or sequence subfragments of the original molecule, such as a human chromosome, would then be assembled from the overlapping F + P sequence determined by computational deduction.

- 5        One option is to transform hybridization signals e.g., scores, into +/- output during the sequence assembly process. In this case, assembly will start with a F + P sequence with a very high score, for example F + P sequence AAAAAATTTTTT. Scores of all four possible overlapping probes AAAAAATTTTTTA , AAAAAATTTTTT , AAAAAATTTTTTC and AAAAAATTTTTTG and three additional probes that are different
- 10 at the beginning (TAAAAATTTTTT , ; CAAAAATTTTTT , ; GAAAAATTTTTT , are compared and three outcomes defined: (i) only the starting probe and only one of the four overlapping probes have scores that are significantly positive relatively to the other six probes, in this case the AAAAAATTTTTT sequence will be extended for one nucleotide to the right; (ii) no one probe except the starting probe has a significantly positive score,
- 15 assembly will stop, e.g., the AAAAAATTTTTT sequence is at the end of the DNA molecule that is sequenced; (iii) more than one significantly positive probe among the overlapped and/or other three probes is found; assembly is stopped because of the error or branching (Drmanac *et al.*, 1989).

- 20        The processes of computational deduction would employ computer programs using existing algorithms (see, e.g., Pevzner, 1989; Drmanac *et al.*, 1991; Labat and Drmanac, 1993; each incorporated herein by reference).

- If, in addition to F + P, F (space 1)P, F (space 2)P, F(space 3)P or F(space 4)P are determined, algorithms will be used to match all data sets to correct potential errors or to solve the situation where there is a branching problem (see, e.g., Drmanac *et al.*,
- 25 1989; Bains *et al.*, 1988; each incorporated herein by reference).

### **EXAMPLE 15**

#### **Conducting Sequencing by Two Step Hybridization**

- Following the certain examples to describe the execution of the sequencing
- 30 methodology contemplated by the inventor. First, the whole chip would be hybridized with mixture of DNA as complex as 100 million of bp (one human chromosome). Guidelines for conducting hybridization can be found in papers such as Drmanac *et al.*

WO 99/09217

PCT/US98/16966

(1990); Khrapko *et al.* (1991); and Broude *et al.* (1994). These articles teach the ranges of hybridization temperatures, buffers and washing steps that are appropriate for use in the initial steps of Format 3 SBH.

The present inventor particularly contemplates that hybridization is to be carried  
5 out for up to several hours in high salt concentrations at a low temperature (-2°C to 5°C) because of a relatively low concentration of target DNA that can be provided. For this purpose, SSC buffer is used instead of sodium phosphate buffer (Drmanac *et al.*, 1990), which precipitates at 10°C. Washing does not have to be extensive (a few minutes) because of the second step, and can be completely eliminated when the hybridization  
10 cycling is used for the sequencing of highly complex DNA samples. The same buffer is used for hybridization and washing steps to be able to continue with the second hybridization step with labeled probes.

After proper washing using a simple robotic device on each array, e.g., a 8 x 8 mm array, one labeled, probe, e.g., a 6-mer, would be added. A 96-tip or 96-pin device  
15 would be used, performing this in 42 operations. Again, a range of discriminatory conditions could be employed, as previously described in the scientific literature.

The present inventor particularly contemplates the use of the following conditions. First, after adding labeled probes and incubating for several minutes only (because of the high concentration of added oligonucleotides) at a low temperature (0-5°C), the  
20 temperature is increased to 3-10°C, depending on F + P length, and the washing buffer is added. At this time, the washing buffer used is one compatible with any ligation reaction (e.g., 100 mM salt concentration range). After adding ligase, the temperature is increased again to 15-37°C to allow fast ligation (less than 30 min) and further discrimination of full match and mismatch hybrids.

25 The use of cationic detergents is also contemplated for use in Format 3 SBH, as described by Pontius & Berg (1991, incorporated herein by reference). These authors describe the use of two simple cationic detergents, dodecyl- and cetyltrimethylammonium bromide (DTAB and CTAB) in DNA renaturation.

DTAB and CTAB are variants of the quaternary amine tetramethylammonium  
30 bromide (TMAB) in which one of the methyl groups is replaced by either a 12-carbon (DTAB) or a 16-carbon (CTAB) alkyl group. TMAB is the bromide salt of the tetramethylammonium ion, a reagent used in nucleic acid renaturation experiments to

WO 99/09217

PCT/US98/16966

decrease the G-C content bias of the melting temperature. DTAB and CTAB are similar in structure to sodium dodecyl sulfate (SDS), with the replacement of the negatively charged sulfate of SDS by a positively charged quaternary amine. While SDS is commonly used in hybridization buffers to reduce nonspecific binding and inhibit  
5 nucleases, it does not greatly affect the rate of renaturation.

When using a ligation process, the enzyme could be added with the labeled probes or after the proper washing step to reduce the background. Although not previously proposed for use in any SBH method, ligase technology is well established within the field of molecular biology. For example, Hood and colleagues described a  
10 ligase-mediated gene detection technique (Landegren *et al.*, 1988), the methodology of which can be readily adapted for use in Format 3 SBH. Wu & Wallace also describe the use of bacteriophage T4 DNA ligase to join two adjacent, short synthetic oligonucleotides. Their oligo ligation reactions were carried out in 50 mM Tris HCl pH 7.6, 10 mM MgCl<sub>2</sub>, 1 mM ATP, 1 mM DTT, and 5% PEG. Ligation reactions were heated to 100°C  
15 for 5-10 min followed by cooling to 0°C prior to the addition of T4 DNA ligase (1 unit; Bethesda Research Laboratory). Most ligation reactions were carried out at 30°C and terminated by heating to 100°C for 5 min.

Final washing appropriate for discriminating detection of hybridized adjacent, or ligated, oligonucleotides of length (F + P), is then performed. This washing step is done  
20 in water for several minutes at 40-60°C to wash out all the non-ligated labeled probes, and all other compounds, to maximally reduce background. Because of the covalently bound labeled oligonucleotides, detection is simplified (it does not have time and low temperature constraints).

Depending on the label used, imaging of the chips is done with different apparatus.  
25 For radioactive labels, phosphor storage screen technology and PhosphorImager as a scanner may be used (Molecular Dynamics, Sunnyvale, CA). Chips are put in a cassette and covered by a phosphorous screen. After 1-4 hours of exposure, the screen is scanned and the image file stored at a computer hard disc. For the detection of fluorescent labels, CCD cameras and epifluorescent or confocal microscopy are used. For the chips  
30 generated directly on the pixels of a CCD camera, detection can be performed as described by Eggers *et al.* (1994, incorporated herein by reference).

WO 99/09217

PCT/US98/16966

Charge-coupled device (CCD) detectors serve as active solid supports that quantitatively detect and image the distribution of labeled target molecules in probe-based assays. These devices use the inherent characteristics of microelectronics that accommodate highly parallel assays, ultrasensitive detection, high throughput, integrated data acquisition and computation. Eggers *et al.* (1994) describe CCDs for use with probe-based assays, such as Format 3 SBH of the present invention, that allow quantitative assessment within seconds due to the high sensitivity and direct coupling employed.

The integrated CCD detection approach enables the detection of molecular binding events on chips. The detector rapidly generates a two-dimensional pattern that uniquely characterizes the sample. In the specific operation of the CCD-based molecular detector, distinct biological probes are immobilized directly on the pixels of a CCD or can be attached to a disposable cover slip placed on the CCD surface. The sample molecules can be labeled with radioisotope, chemiluminescent or fluorescent tags.

Upon exposure of the sample to the CCD-based probe array, photons or radioisotope decay products are emitted at the pixel locations where the sample has bound, in the case of Format 3, to two complementary probes. In turn, electron-hole pairs are generated in the silicon when the charged particles, or radiation from the labeled sample, are incident on the CCD gates. Electrons are then collected beneath adjacent CCD gates and sequentially read out on a display module. The number of photoelectrons generated at each pixel is directly proportional to the number of molecular binding events in such proximity. Consequently, molecular binding can be quantitatively determined (Eggers *et al.*, 1994).

By placing the imaging array in proximity to the sample, the collection efficiency is improved by a factor of at least 10 over lens-based techniques such as those found in conventional CCD cameras. That is, the sample (emitter) is in near contact with the detector (imaging array), and this eliminates conventional imaging optics such as lenses and mirrors.

When radioisotopes are attached as reporter groups to the target molecules, energetic particles are detected. Several reporter groups that emit particles of varying energies have been successfully utilized with the micro-fabricated detectors, including  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^{35}\text{S}$ ,  $^{14}\text{C}$  and  $^{125}\text{I}$ . The higher energy particles, such as from  $^{32}\text{P}$ , provide the highest

WO 99/09217

PCT/US98/16966

molecular detection sensitivity, whereas the lower energy particles, such as from  $^{35}\text{S}$ , provide better resolution. Hence the choice of the radioisotope reporter can be tailored as required. Once the particular radioisotope label is selected, the detection performance can be predicted by calculating the signal-to-noise ration (SNR), as described by Eggers *et al.* (1994).

An alternative luminescent detection procedure involves the use of fluorescent or chemiluminescent reporter groups attached to the target molecules. The fluorescent labels can be attached covalently or through interaction. Fluorescent dyes, such as ethidium bromide, with intense absorption bands in the near UV (300-350 nm) range and principal emission bands in the visible (500-650 nm) range, are most suited for the CCD devices employed since the quantum efficiency is several orders of magnitude lower at the excitation wavelength than at the fluorescent signal wavelength.

From the perspective of detecting luminescence, the polysilicon CCD gates have the built-in capacity to filter away the contribution of incident light in the UV range, yet are very sensitive to the visible luminescence generated by the fluorescent reporter groups. Such inherently large discrimination against UV excitation enables large SNRs (greater than 100) to be achieved by the CCDs as formulated in the incorporated paper by Eggers *et al.* (1994).

For probe immobilization on the detector, hybridization matrices may be produced on inexpensive  $\text{SiO}_2$  wafers, which are subsequently placed on the surface of the CCD following hybridization and drying. This format is economically efficient since the hybridization of the DNA is conducted on inexpensive disposable  $\text{SiO}_2$  wafers, thus allowing reuse of the more expensive CCD detector. Alternatively, the probes can be immobilized directly on the CCD to create a dedicated probe matrix.

To immobilize probes upon the  $\text{SiO}_2$  coating, a uniform epoxide layer is linked to the film surface, employing an epoxy-silane reagent and standard  $\text{SiO}_2$  modification chemistry. Amine-modified oligonucleotide probes are then linked to the  $\text{SiO}_2$  surface by means of secondary amine formation with the epoxide ring. The resulting linkage provides 17 rotatable bonds of separation between the 3' base of the oligonucleotide and the  $\text{SiO}_2$  surface. To ensure complete amine deprotonation and to minimize secondary structure formation during coupling, the reaction is performed in 0.1 M KOH and incubated at 37°C for 6 hours.



WO 99/09217

PCT/US98/16966

In Format 3 SBH in general, signals are scored per each of billion points. It would not be necessary to hybridize all arrays, e.g., 4000 5 x 5 mm, at a time and the successive use of smaller number of arrays is possible.

Cycling hybridizations are one possible method for increasing the hybridization  
5 signal. In one cycle, most of the fixed probes will hybridize with DNA fragments with  
tail sequences non-complementary for labeled probes. By increasing the temperature,  
those hybrids will be melted. In the next cycle, some of them (~0.1%) will hybridize  
with an appropriate DNA fragment and additional labeled probes will be ligated. In this  
case, there occurs a discriminative melting of DNA hybrids with mismatches for both  
10 probe sets simultaneously.

In the cycle hybridization, all components are added before the cycling starts, at  
the 37°C for T4, or a higher temperature for a thermostable ligase. Then the temperature  
is decreased to 15-37°C and the chip is incubated for up to 10 minutes, and then the  
temperature is increased to 37°C or higher for a few minutes and then again reduced.  
15 Cycles can be repeated up to 10 times. In one variant, an optimal higher temperature  
(10-50°C) can be used without cycling and longer ligation reaction can be performed (1-3  
hours).

The procedure described herein allows complex chip manufacturing using standard  
synthesis and precise spotting of oligonucleotides because a relatively small number of  
20 oligonucleotides are necessary. For example, if all 7-mer oligos are synthesized (16384  
probes), lists of 256 million 14-mers can be determined.

One important variant of the invented method is to use more than one differently  
labeled probe per base array. This can be executed with two purposes in mind;  
multiplexing to reduce number of separately hybridized arrays; or to determine a list of  
25 even longer oligosequences such as 3 x 6 or 3 x 7. In this case, if two labels are used,  
the specificity of the 3 consecutive oligonucleotides can be almost absolute because  
positive sites must have enough signals of both labels.

A further and additional variant is to use chips containing BxNy probes with y  
being from 1 to 4. Those chips allow sequence reading in different frames. This can  
30 also be achieved by using appropriate sets of labeled probes or both F and P probes could  
have some unspecified end positions (i.e., some element of terminal degeneracy).  
Universal bases may also be employed as part of a linker to join the probes of defined

WO 99/09217

PCT/US98/16966

sequence to the solid support. This makes the probe more available to hybridization and makes the construct more stable. If a probe has 5 bases, one may, e.g., use 3 universal bases as a linker.

## 5 **EXAMPLE 16**

### **Determining Sequence from Hybridization Data**

Sequence assembly may be interrupted where ever a given overlapping (N-1) mer is duplicated two or more times. Then either of the two N-mers differing in the last nucleotide may be used in extending the sequence. This branching point limits

10 unambiguous assembly of sequence.

Reassembling the sequence of known oligonucleotides that hybridize to the target nucleic acid to generate the complete sequence of the target nucleic acid may not be accomplished in some cases. This is because some information may be lost if the target nucleic acid is not in fragments of appropriate size in relation to the size of

15 oligonucleotide that is used for hybridizing. The quantity of information lost is proportional to the length of a target being sequenced. However, if sufficiently short targets are used, their sequence may be unambiguously determined.

The probable frequency of duplicated sequences that would interfere with sequence assembly which is distributed along a certain length of DNA may be calculated. This

20 derivation requires the introduction of the definition of a parameter having to do with sequence organization: the sequence subfragment (SF). A sequence subfragment results if any part of the sequence of a target nucleic acid starts and ends with an (N-1)mer that is repeated two or more times within the target sequence. Thus, subfragments are sequences generated between two points of branching in the process of assembly of the

25 sequences in the method of the invention. The sum of all subfragments is longer than the actual target nucleic acid because of overlapping short ends. Generally, subfragments may not be assembled in a linear order without additional information since they have shared (N-1)mers at their ends and starts. Different numbers of subfragments are obtained for each nucleic acid target depending on the number of its repeated (N-1) mers.

30 The number depends on the value of N-1 and the length of the target.

Probability calculations can estimate the interrelationship of the two factors. If the ordering of positive N-mers is accomplished by using overlapping sequences of length

WO 99/09217

PCT/US98/16966

N-1 or at an average distance of  $A_0$ , the N-1 of a fragment  $L_f$  bases long is given by equation one:

$$N_{fr} = 1 + A_0 \times K \times P(K, L_f)$$

Where  $K$  greater than or  $= 2$ , and  $P(K, L_f)$  represents the probability of an  
 5 N-mer occurring  $K$ -times on a fragment  $L_f$  base long. Also, a computer program that is able to form subfragments from the content of N-mers for any given sequence is described below in Example 18.

The number of subfragments increases with the increase of lengths of fragments for a given length of probe. Obtained subfragments may not be uniquely ordered among  
 10 themselves. Although not complete, this information is very useful for comparative sequence analysis and the recognition of functional sequence characteristics. This type of information may be called partial sequence. Another way of obtaining partial sequence is the use of only a subset of oligonucleotide probes of a given length.

There may be relatively good agreement between predicted sequence according to  
 15 theory and a computer simulation for a random DNA sequence. For instance, for  $N-1 = 7$ , [using an 8-mer or groups of sixteen 10-mers of type  $5' (A,T,C,G) B_8 (A,T,C,G) 3'$ ] a target nucleic acid of 200 bases will have an average of three subfragments. However, because of the dispersion around the mean, a library of target nucleic acid should have inserts of 500 bp so that less than 1 in 2000 targets have more than three subfragments.  
 20 Thus, in an ideal case of sequence determination of a long nucleic acid of random sequence, a representative library with sufficiently short inserts of target nucleic acid may be used. For such inserts, it is possible to reconstruct the individual target by the method of the invention. The entire sequence of a large nucleic acid is then obtained by overlapping of the defined individual insert sequences.

25 To reduce the need for very short fragments, e.g. 50 bases for 8-mer probes. The information contained in the overlapped fragments present in every random DNA fragmentation process like cloning, or random PCR is used. It is also possible to use pools of short physical nucleic acid fragments. Using 8-mers or 11-mers like  $5' (A, T, C, G) N_8 (A, T, C, G) 3'$  for sequencing 1 megabase, instead of needing 20,000 50 bp  
 30 fragments only 2,100 samples are sufficient. This number consists of 700 random 7 kb clones (basic library), 1250 pools of 20 clones of 500 bp (subfragments ordering library)

WO 99/09217

PCT/US98/16966

and 150 clones from jumping (or similar) library. The developed algorithm (see Example 18) regenerates sequence using hybridization data of these described samples.

### **EXAMPLE 17**

#### **5 Algorithm**

This example describes an algorithm for generation of a long sequence written in a four letter alphabet from constituent k-tuple words in a minimal number of separate, randomly defined fragments of a starting nucleic acid sequence where K is the length of an oligonucleotide probe. The algorithm is primarily intended for use in the sequencing  
10 by hybridization (SBH) process. The algorithm is based on subfragments (SF), informative fragments (IF) and the possibility of using pools of physical nucleic sequences for defining informative fragments.

As described, subfragments may be caused by branch points in the assembly process resulting from the repetition of a K-1 oligomer sequence in a target nucleic acid.  
15 Subfragments are sequence fragments found between any two repetitive words of the length K-1 that occur in a sequence. Multiple occurrences of K-1 words are the cause of interruption of ordering the overlap of K-words in the process of sequence generation. Interruption leads to a sequence remaining in the form of subfragments. Thus, the unambiguous segments between branching points whose order is not uniquely determined  
20 are called sequence subfragments.

Informative fragments are defined as fragments of a sequence that are determined by the nearest ends of overlapped physical sequence fragments.

A certain number of physical fragments may be pooled without losing the possibility of defining informative fragments. The total length of randomly pooled  
25 fragments depends on the length of k-tuples that are used in the sequencing process.

The algorithm consists of two main units. The first part is used for generation of subfragments from the set of k-tuples contained in a sequence. Subfragments may be generated within the coding region of physical nucleic acid sequence of certain sizes, or within the informative fragments defined within long nucleic acid sequences. Both types  
30 of fragments are members of the basic library. This algorithm does not describe the determination of the content of the k-tuples of the informative fragments of the basic

WO 99/09217

PCT/US98/16966

library, i.e. the step of preparation of informative fragments to be used in the sequence generation process.

The second part of the algorithm determines the linear order of obtained subfragments with the purpose of regenerating the complete sequence of the nucleic acid fragments of the basic library. For this purpose a second, ordering library is used, made of randomly pooled fragments of the starting sequence. The algorithm does not include the step of combining sequences of basic fragments to regenerate an entire, megabase plus sequence. This may be accomplished using the link-up of fragments of the basic library which is a prerequisite for informative fragment generation. Alternatively, it may be accomplished after generation of sequences of fragments of the basic library by this algorithm, using search for their overlap, based on the presence of common end-sequences.

The algorithm requires neither knowledge of the number of appearances of a given k-tuple in a nucleic acid sequence of the basic and ordering libraries, nor does it require the information of which k-tuple words are present on the ends of a fragment. The algorithm operates with the mixed content of k-tuples of various length. The concept of the algorithm enables operations with the k-tuple sets that contain false positive and false negative k-tuples. Only in specific cases does the content of the false k-tuples primarily influence the completeness and correctness of the generated sequence. The algorithm may be used for optimization of parameters in simulation experiments, as well as for sequence generation in the actual SBH experiments e.g. generation of the genomic DNA sequence. In optimization of parameters, the choice of the oligonucleotide probes (k-tuples) for practical and convenient fragments and/or the choice of the optimal lengths and the number of fragments for the defined probes are especially important.

This part of the algorithm has a central role in the process of the generation of the sequence from the content of k-tuples. It is based on the unique ordering of k-tuples by means of maximal overlap. The main obstacles in sequence generation are specific repeated sequences and false positive and/or negative k-tuples. The aim of this part of the algorithm is to obtain the minimal number of the longest possible subfragments, with correct sequence. This part of the algorithm consists of one basic, and several control steps. A two-stage process is necessary since certain information can be used only after generation of all primary subfragments.

WO 99/09217

PCT/US98/16966

The main problem of sequence generation is obtaining a repeated sequence from word contents that by definition do not carry information on the number of occurrences of the particular k-tuples. The concept of the entire algorithm depends on the basis on which this problem is solved. In principle, there are two opposite approaches: 1) repeated sequences may be obtained at the beginning, in the process of generation of pSFs, or 2) repeated sequences can be obtained later, in the process of the final ordering of the subfragments. In the first case, pSFs contain an excess of sequences and in the second case, they contain a deficit of sequences. The first approach requires elimination of the excess sequences generated, and the second requires permitting multiple use of some of the subfragments in the process of the final assembling of the sequence.

The difference in the two approaches in the degree of strictness of the rule of unique overlap of k-tuples. The less severe rule is: k-tuple X is unambiguously maximally overlapped with k-tuple Y if and only if, the rightmost k-1 end of k-tuple X is present only on the leftmost end of k-tuple Y. This rule allows the generation of repetitive sequences and the formation of surplus sequences.

A stricter rule which is used in the second approach has an addition caveat: k-tuple X is unambiguously maximally overlapped with k-tuple Y if and only if, the rightmost K-1 end of k-tuple X is present only on the leftmost end of k-tuple Y and if the leftmost K-1 end of k-tuple Y is not present on the rightmost end of any other k-tuple. The algorithm based on the stricter rule is simpler, and is described herein.

The process of elongation of a given subfragment is stopped when the right k-1 end of the last k-tuple included is not present on the left end of any k-tuple or is present on two or more k-tuples. If it is present on only one k-tuple the second part of the rule is tested. If in addition there is a k-tuple which differs from the previously included one, the assembly of the given subfragment is terminated only on the first leftmost position. If this additional k-tuple does not exist, the conditions are met for unique k-1 overlap and a given subfragment is extended to the right by one element.

Beside the basic rule, a supplementary one is used to allow the usage of k-tuples of different lengths. The maximal overlap is the length of k-1 of the shorter k-tuple of the overlapping pair. Generation of the pSFs is performed starting from the first k-tuple from the file in which k-tuples are displayed randomly and independently from their order in a nucleic acid sequence. Thus, the first k-tuple in the file is not necessarily on the

WO 99/09217

PCT/US98/16966

beginning of the sequence, nor on the start of the particular subfragment. The process of subfragment generation is performed by ordering the k-tuples by means of unique overlap, which is defined by the described rule. Each used k-tuple is erased from the file. At the point when there are no further k-tuples unambiguously overlapping with the last one included, the building of subfragment is terminated and the buildup of another pSF is started. Since generation of a majority of subfragments does not begin from their actual starts, the formed pSF are added to the k-tuple file and are considered as a longer k-tuple. Another possibility is to form subfragments going in both directions from the starting k-tuple. The process ends when further overlap, i.e. the extension of any of the subfragments, is not possible.

The pSFs can be divided in three groups: 1) Subfragments of the maximal length and correct sequence in cases of exact k-tuple set; 2) short subfragments, formed due to the used of the maximal and unambiguous overlap rule on the incomplete set, and/or the set with some false positive k-tuples; and 3) pSFs of an incorrect sequence. The incompleteness of the set in 2) is caused by false negative results of a hybridization experiment, as well as by using an incorrect set of k-tuples. These are formed due to the false positive and false negative k-tuples and can be : a) misconnected subfragments; b) subfragments with the wrong end; and c) false positive k-tuples which appears as false minimal subfragments.

Considering false positive k-tuples, there is the possibility for the presence of a k-tuple containing more than one wrong base or containing one wrong base somewhere in the middle, as well as the possibility for a k-tuple with a wrong base on the end. Generation of short, erroneous or misconnected subfragments is caused by the latter k-tuples. The k-tuples of the former two kinds represent wrong pSFs with length equal to k-tuple length.

In the case of one false negative k-tuple, pSFs are generated because of the impossibility of maximal overlapping. In the case of the presence of one false positive k-tuple with the wrong base on its leftmost or rightmost end, pSFs are generated because of the impossibility of unambiguous overlapping. When both false positive and false negative k-tuples with a common k-1 sequence are present in the file, pSFs are generated, and one of these pSFs contains the wrong k-tuple at the relevant end.

WO 99/09217

PCT/US98/16966

The process of correcting subfragments with errors in sequence and the linking of unambiguously connected pSF is performed after subfragment generation and in the process of subfragment ordering. The first step which consists of cutting the misconnected pSFs and obtaining the final subfragments by unambiguous connection of pSFs is described below.

There are two approaches for the formation of misconnected subfragments. In the first a mistake occurs when an erroneous k-tuple appears on the points of assembly of the repeated sequences of lengths k-1. In the second, the repeated sequences are shorter than k-1. These situations can occur in two variants each. In the first variant, one of the repeated sequences represents the end of a fragment. In the second variant, the repeated sequence occurs at any position within the fragment. For the first possibility, the absence of some k-tuples from the file (false negatives) is required to generate a misconnection. The second possibility requires the presence of both false negative and false positive k-tuples in the file. Considering the repetitions of k-1 sequence, the lack of only one k-tuple is sufficient when either end is repeated internally. The lack of two is needed for strictly internal repetition. The reason is that the end of a sequence can be considered informatically as an endless linear array of false negative k-tuples. From the "smaller than k-1 case", only the repeated sequence of the length of k-2, which requires two or three specific erroneous k-tuples, will be considered. It is very likely that these will be the only cases which will be detected in a real experiment, the others being much less frequent.

Recognition of the misconnected subfragments is more strictly defined when a repeated sequence does not appear at the end of the fragment. In this situation, one can detect further two subfragments, one of which contains on its leftmost, and the other on its rightmost end k-2 sequences which are also present in the misconnected subfragment. When the repeated sequence is on the end of the fragment, there is only one subfragment which contains k-2 sequence causing the mistake in subfragment formation on its leftmost or rightmost end.

The removal of misconnected subfragments by their cutting is performed according to the common rule: If the leftmost or rightmost sequence of the length of k-2 of any subfragments is present in any other subfragment, the subfragment is to be cut into two subfragments, each of them containing k-2 sequence. This rule does not cover rarer



WO 99/09217

PCT/US98/16966

situations of a repeated end when there are more than one false negative k-tuple on the point of repeated k-1 sequence. Misconnected subfragments of this kind can be recognized by using the information from the overlapped fragments, or informative fragments of both the basic and ordering libraries. In addition, the misconnected  
5 subfragment will remain when two or more false negative k-tuples occur on both positions which contain the identical k-1 sequence. This is a very rare situation since it requires at least 4 specific false k-tuples. An additional rule can be introduced to cut these subfragments on sequences of length k if the given sequence can be obtained by combination of sequences shorter than k-2 from the end of one subfragment and the start  
10 of another.

By strict application of the described rule, some completeness is lost to ensure the accuracy of the output. Some of the subfragments will be cut although they are not misconnected since they fit into the pattern of a misconnected subfragment. There are several situations of this kind. For example, a fragment, beside at least two identical k-1  
15 sequences, contains any k-2 sequence from k-1 or a fragment contains k-2 sequence repeated at least twice and at least one false negative k-tuple containing given k-2 sequence in the middle, etc.

The aim of this part of the algorithm is to reduce the number of pSFs to a minimal number of longer subfragments with correct sequence. The generation of unique longer  
20 subfragments or a complete sequence is possible in two situations. The first situation concerns the specific order of repeated k-1 words. There are cases in which some or all maximally extended pSFs (the first group of pSFs) can be uniquely ordered. For example, in fragment S-R1-a-R2-b-R1-c-R2-E where S and E are the start and end of a fragment, a, b, and c are different sequences specific to respective subfragments and R1  
25 and R2 are two k-1 sequences that are tandemly repeated, five subfragments are generated (S-R1, R1-a-R2, R2-b-R1, R1-c-R2, and R-E). They may be ordered in two ways; the original sequence above or S-R1- c-R-b-R1-a-R-E. In contrast, in a fragment with the same number and types of repeated sequences but ordered differently, i.e. S-R1-a-R1-b-R-c-R-E, there is no other sequence which includes all subfragments.  
30 Examples of this type can be recognized only after the process of generation of pSFs. They represent the necessity for two steps in the process of pSF generation. The second situation of generation of false short subfragments on positions of nonrepeated k-1

WO 99/09217

PCT/US98/16966

sequences when the files contain false negative and /or positive k-tuples is more important.

The solution for both pSF groups consists of two parts. First, the false positive k-tuples appearing as the nonexisting minimal subfragments are eliminated. All k-tuple  
5 subfragments of length k which do not have an overlap on either end, of the length of longer than k-a on one end and longer than k-b on the other end, are eliminated to enable formation of the maximal number of connections. In our experiments, the values for a and b of 2 and 3, respectively, appeared to be adequate to eliminate a sufficient number of false positive k-tuples.

10 The merging of subfragments that can be uniquely connected is accomplished in the second step. The rule for connection is: two subfragments may be unambiguously connected if, and only if, the overlapping sequence at the relevant end or start of two subfragments is not present at the start and/or end of any other subfragment.

The exception is if one subfragment from the considered pair has the identical  
15 beginning and end. In that case connection is permitted, even if there is another subfragment with the same end present in the file. The main problem here is the precise definition of overlapping sequence. The connection is not permitted if the overlapping sequence unique for only one pair of subfragments is shorter than k-2, of it is k-2 or longer but an additional subfragment exists with the overlapping sequence of any length  
20 longer than k-4. Also, both the canonical ends of pSFs and the ends after omitting one (or few) last bases are considered as the overlapping sequences.

After this step some false positive k-tuples (as minimal subfragments) and some subfragments with a wrong end may survive. In addition, in very rare occasions where a certain number of some specific false k-tuples are simultaneously present, an erroneous  
25 connection may take place. These cases will be detected and solved in the subfragment ordering process, and in the additional control steps along with the handling of uncut "misconnected" subfragments.

The short subfragments that are obtained are of two kinds. In the common case, these subfragments may be unambiguously connected among themselves because of the  
30 distribution of repeated k-1 sequences. This may be done after the process of generation of pSFs and is a good example of the necessity for two steps in the process of pSF generation. In the case of using the file containing false positive and/or false negative

WO 99/09217

PCT/US98/16966

k-tuples, short pSFs are obtained on the sites of nonrepeated k-1 sequences. Considering false positive k-tuples, a k-tuple may contain more than one wrong base (or containing one wrong base somewhere in the middle), as well as k-tuple on the end. Generation of short and erroneous (or misconnected) subfragments is caused by the latter k-tuples. The

5 k-tuples of the former kind represent wrong pSFs with length equal to k- tuple length.

The aim of merging pSF part of the algorithm is the reduction of the number of pSFs to the minimal number of longer subfragments with the correct sequence. All k-tuple subfragments that do not have an overlap on either end, of the length of longer than k-a on one, and longer than k-b on the other end, are eliminated to enable the

10 maximal number of connections. In this way, the majority of false positive k-tuples are discarded. The rule for connection is: two subfragments can be unambiguously connected if, and only if the overlapping sequence of the relevant end or start of two subfragments is not present on the start and/or end of any other subfragment. The exception is a subfragment with the identical beginning and end. In that case connection

15 is permitted, provided that there is another subfragment with the same end present in the file. The main problem here is of precise definition of overlapping sequence. The presence of at least two specific false negative k- tuples on the points of repetition of k-1 or k-2 sequences, as well as combining of the false positive and false negative k-tuples may destroy or "mask" some overlapping sequences and can produce an unambiguous,

20 but wrong connection of pSFs. To prevent this, completeness must be sacrificed on account of exactness: the connection is not permitted on the end-sequences shorter than k-2, and in the presence of an extra overlapping sequence longer than k-4. The overlapping sequences are defined from the end of the pSFs, or omitting one, or few last bases.

25 In the very rare situations, with the presence of a certain number of some specific false positive and false negative k-tuples, some subfragments with the wrong end can survive, some false positive k-tuples (as minimal subfragments) can remain, or the erroneous connection can take place. These cases are detected and solved in the subfragments ordering process, and in the additional control steps along with the handling

30 of uncut, misconnected subfragments.

The process of ordering of subfragments is similar to the process of their generation. If one considers subfragments as longer k-tuples, ordering is performed by

WO 99/09217

PCT/US98/16966

their unambiguous connection via overlapping ends. The informational basis for unambiguous connection is the division of subfragments generated in fragments of the basic library into groups representing segments of those fragments. The method is analogous to the biochemical solution of this problem based on hybridization with longer  
5 oligonucleotides with relevant connecting sequence. The connecting sequences are generated as subfragments using the k-tuple sets of the appropriate segments of basic library fragments. Relevant segments are defined by the fragments of the ordering library that overlap with the respective fragments of the basic library. The shortest segments are informative fragments of the ordering library. The longer ones are several neighboring  
10 informative fragments or total overlapping portions of fragments corresponding of the ordering and basic libraries. In order to decrease the number of separate samples, fragments of the ordering library are randomly pooled, and the unique k- tuple content is determined.

By using the large number of fragments in the ordering library very short  
15 segments are generated, thus reducing the chance of the multiple appearance of the k-1 sequences which are the reasons for generation of the subfragments. Furthermore, longer segments, consisting of the various regions of the given fragment of the basic library, do not contain some of the repeated k-1 sequences. In every segment a connecting sequence (a connecting subfragment) is generated for a certain pair of the subfragments from the  
20 given fragment. The process of ordering consists of three steps: (1) generation of the k-tuple contents of each segment; (2) generation of subfragments in each segment; and (3) connection of the subfragments of the segments. Primary segments are defined as significant intersections and differences of k-tuple contents of a given fragment of the basic library with the k-tuple contents of the pools of the ordering library. Secondary  
25 (shorter) segments are defined as intersections and differences of the k-tuple contents of the primary segments.

There is a problem of accumulating both false positive and negative k-tuples in both the differences and intersections. The false negative k-tuples from starting sequences accumulate in the intersections (overlapping parts), as well as false positive k-tuples  
30 occurring randomly in both sequences, but not in the relevant overlapping region. On the other hand, the majority of false positives from either of the starting sequences is not taken up into intersections. This is an example of the reduction of experimental errors

WO 99/09217

PCT/US98/16966

from individual fragments by using information from fragments overlapping with them. The false k- tuples accumulate in the differences for another reason. The set of false negatives from the original sequences are enlarged for false positives from intersections and the set of false positives for those k-tuples which are not included in the intersection  
5 by error, i.e. are false negative in the intersection. If the starting sequences contain 10% false negative data, the primary and secondary intersections will contain 19% and 28% false negative k- tuples, respectively. On the other hand, a mathematical expectation of 77 false positives may be predicted if the basic fragment and the pools have lengths of 500 bp and 10,000 bp, respectively. However, there is a possibility of recovering most  
10 of the "lost" k-tuples and of eliminating most of the false positive k-tuples.

First, one has to determine a basic content of the k-tuples for a given segment as the intersection of a given pair of the k-tuple contents. This is followed by including all k- tuples of the starting k-tuple contents in the intersection, which contain at one end k-1 and at the other end k-+ sequences which occur at the ends of two k-tuples of the basic  
15 set. This is done before generation of the differences thus preventing the accumulation of false positives in that process. Following that, the same type of enlargement of k-tuple set is applied to differences with the distinction that the borrowing is from the intersections. All borrowed k-tuples are eliminated from the intersection files as false positives.

20 The intersection, i.e. a set of common k-tuples, is defined for each pair (a basic fragment) X (a pool of ordering library). If the number of k-tuples in the set is significant it is enlarged with the false negatives according to the described rule. The primary difference set is obtained by subtracting from a given basic fragment the obtained intersection set. The false negative k-tuples are appended to the difference set by  
25 borrowing from the intersection set according to the described rule and, at the same time, removed from the intersection set as false positive k-tuples. When the basic fragment is longer than the pooled fragments, this difference can represent the two separate segments which somewhat reduces its utility in further steps. The primary segments are all generated intersections and differences of pairs (a basic fragment) X (a pool of ordering  
30 library) containing the significant number of k-tuples. K-tuple sets of secondary segments are obtained by comparison of k-tuple sets of all possible pairs of primary segments. The two differences are defined from each pair which produces the intersection with the

WO 99/09217

PCT/US98/16966

significant number of k-tuples. The majority of available information from overlapped fragments is recovered in this step so that there is little to be gained from the third round of forming intersections, and differences.

(2) Generation of the subfragments of the segments is performed identically as  
5 described for the fragments of the basic library.

(3) The method of connection of subfragments consists of sequentially  
determining the correctly linked pairs of subfragments among the subfragments from a  
given basic library fragment which have some overlapped ends. In the case of 4 relevant  
subfragments, two of which contain the same beginning and two having the same end,  
10 there are 4 different pairs of subfragments that can be connected. In general 2 are correct  
and 2 are wrong. To find correct ones, the presence of the connecting sequences of each  
pair is tested in the subfragments generated from all primary and secondary segments for  
a given basic fragment. The length and the position of the connecting sequence are  
chosen to avoid interference with sequences which occur by chance. They are  $k+2$  or  
15 longer, and include at least one element 2 beside overlapping sequence in both  
subfragments of a given pair. The connection is permitted only if the two connecting  
sequences are found and the remaining two do not exist. The two linked subfragments  
replace former subfragments in the file and the process is cyclically repeated.

Repeated sequences are generated in this step. This means that some subfragments  
20 are included in linked subfragments more than once. They will be recognized by finding  
the relevant connecting sequence which engages one subfragment in connection with two  
different subfragments.

The recognition of misconnected subfragments generated in the processes of  
building pSFs and merging pSFs into longer subfragments is based on testing whether the  
25 sequences of subfragments from a given basic fragment exist in the sequences of  
subfragments generated in the segments for the fragment. The sequences from an  
incorrectly connected position will not be found indicating the misconnected  
subfragments.

Beside the described three steps in ordering of subfragments some additional  
30 control steps or steps applicable to specific sequences will be necessary for the generation  
of more complete sequence without mistakes.

WO 99/09217

PCT/US98/16966

The determination of which subfragment belongs to which segment is performed by comparison of contents of k-tuples in segments and subfragments. Because of the errors in the k-tuple contents (due to the primary error in pools and statistical errors due to the frequency of occurrences of k-tuples) the exact partitioning of subfragments is impossible.

- 5 Thus, instead of "all or none" partition, the chance of coming from the given segment ( $P(sf,s)$ ) is determined for each subfragment. This possibility is the function of the lengths of k-tuples, the lengths of subfragments, the lengths of fragments of ordering library, the size of the pool, and of the percentage of false k-tuples in the file:

$$P(sf,s) = (C_k - F) / L_{sf},$$

- 10 where  $L_{sf}$  is the length of subfragment,  $C_k$  is the number of common k-tuples for a given subfragment/segment pair, and  $F$  is the parameter that includes relations between lengths of k-tuples, fragments of basic library, the size of the pool, and the error percentage.

- Subfragments attributed to a particular segment are treated as redundant short pSFs and are submitted to a process of unambiguous connection. The definition of
- 15 unambiguous connection is slightly different in this case, since it is based on a probability that subfragments with overlapping end(s) belong to the segment considered. Besides, the accuracy of unambiguous connection is controlled by following the connection of these subfragments in other segments. After the connection in different segments, all of the obtained subfragments are merged together, shorter subfragments included within longer
- 20 ones are eliminated, and the remaining ones are submitted to the ordinary connecting process. If the sequence is not regenerated completely, the process of partition and connection of subfragments is repeated with the same or less severe criterions of probability of belonging to the particular segment, followed by unambiguous connection.

- Using severe criteria for defining unambiguous overlap, some information is not
- 25 used. Instead of a complete sequence, several subfragments that define a number of possibilities for a given fragment are obtained. Using less severe criteria an accurate and complete sequence is generated. In a certain number of situations, e.g. an erroneous connection, it is possible to generate a complete, but an incorrect sequence, or to generate "monster" subfragments with no connection among them. Thus, for each fragment of the
- 30 basic library one obtains: a) several possible solutions where one is correct and b) the most probable correct solution. Also, in a very small number of cases, due to the mistake in the subfragment generation process or due to the specific ratio of the

WO 99/09217

PCT/US98/16966

probabilities of belonging, no unambiguous solution is generated or one, the most probable solution. These cases remain as incomplete sequences, or the unambiguous solution is obtained by comparing these data with other, overlapped fragments of basic library.

5       The described algorithm was tested on a randomly generated, 50 kb sequence, containing 40% GC to simulate the GC content of the human genome. In the middle part of this sequence were inserted various AlI, and some other repetitive sequences, of a total length of about 4 kb. To simulate an in vitro SBH experiment, the following operations were performed to prepare appropriate data.

10       - Positions of sixty 5 kb overlapping "clones" were randomly defined, to simulate preparation of a basic library:

      - Positions of one thousand 500 bp "clones" were randomly determined to simulate making the ordering library. These fragments were extracted from the sequence.

      Random pools of 20 fragments were made, and k-tuple sets of pools were determined and  
15       stored on the hard disk. These data are used in the subfragment ordering phase: For the same density of clones 4 million clones in basic library and 3 million clones in ordering library are used for the entire human genome. The total number of 7 million clones is several fold smaller than the number of clones a few kb long for random cloning of almost all of genomic DNA and sequencing by a gel-based method.

20       From the data on the starts and ends of 5 kb fragments, 117 "informative fragments" were determined to be in the sequence. This was followed by determination of sets of overlapping k-tuples of which the single "informative fragment" consist. Only the subset of k-tuples matching a predetermined list were used. The list contained 65% 8-mers, 30% 9-mers, and 5% 10-12-mers. Processes of generation and the ordering of  
25       subfragments were performed on these data.

      The testing of the algorithm was performed on the simulated data in two experiments. The sequence of 50 informative fragments was regenerated with the 100% correct data set (over 20,000 bp), and 26 informative fragments (about 10,000 bp) with 10% false k-tuples (5% positive and 5% negative ones).

30       In the first experiment, all subfragments were correct and in only one out of 50 informative fragments the sequence was not completely regenerated but remained in the form of 5 subfragments. The analysis of positions of overlapped fragments of ordering



WO 99/09217

PCT/US98/16966

library has shown that they lack the information for the unique ordering of the 5 subfragments. The subfragments may be connected in two ways based on overlapping ends, 1-2-3-4-5 and 1-4-3-2-5. The only difference is the exchange of positions of subfragments 2 and 4. Since subfragments 2, 3, and 4 are relatively short (total of about 5 100 bp), the relatively greater chance existed, and occurred in this case, that none of the fragments of ordering library started or ended in the subfragment 3 region.

To simulate real sequencing, some false ("hybridization") data was included as input in a number of experiments. In oligomer hybridization experiments, under proposed conditions, the only situation producing unreliable data is the end mismatch 10 versus full match hybridization. Therefore, in simulation only those k-tuples differing in a single element on either end from the real one were considered to be false positives. These "false" sets are made as follows. On the original set of a k-tuples of the informative fragment, a subset of 5% false positive k-tuples are added. False positive k-tuples are made by randomly picking a k-tuple from the set, copying it and altering a 15 nucleotide on its beginning or end. This is followed by subtraction of a subset of 5% randomly chosen k-tuples. In this way the statistically expected number of the most complicated cases is generated in which the correct k-tuple is replaced with a k-tuple with the wrong base on the end.

Production of k-tuple sets as described leads to up to 10% of false data. This 20 value varies from case to case, due to the randomness of choice of k-tuples to be copied, altered, and erased. Nevertheless, this percentage 3-4 times exceeds the amount of unreliable data in real hybridization experiments. The introduced error of 10% leads to the two fold increase in the number of subfragments both in fragments of basic library (basic library informative fragments) and in segments. About 10% of the final 25 subfragments have a wrong base at the end as expected for the k-tuple set which contains false positives (see generation of primary subfragments). Neither the cases of misconnection of subfragments nor subfragments with the wrong sequence were observed. In 4 informative fragments out of 26 examined in the ordering process the complete sequence was not regenerated. In all 4 cases the sequence was obtained in the form of 30 several longer subfragments and several shorter subfragments contained in the same segment. This result shows that the algorithmic principles allow working with a large percentage of false data.

WO 99/09217

PCT/US98/16966

The success of the generation of the sequence from its k-tuple content may be described in terms of completeness and accuracy. In the process of generation, two particular situations can be defined: 1) Some part of the information is missing in the generated sequence, but one knows where the ambiguities are and to which type they belong, and 2) the regenerated sequence that is obtained does not match the sequence from which the k-tuple content is generated, but the mistake can not be detected. Assuming the algorithm is developed to its theoretical limits, as in the use of the exact k-tuple sets, only the first situation can take place. There the incompleteness results in a certain number of subfragments that may not be ordered unambiguously and the problem of determination of the exact length of monotonous sequences, i.e. the number of perfect tandem repeats.

With false k-tuples, incorrect sequence may be generated. The reason for mistakes does not lie in the shortcomings of the algorithm, but in the fact that a given content of k-tuples unambiguously represents the sequence that differs from the original one. One may define three classes of error, depending on the kind of the false k-tuples present in the file. False negative k-tuples (which are not accompanied with the false positives) produce "deletions". False positive k-tuples are producing "elongations (unequal crossing over)". False positives accompanied with false negatives are the reason for generation of "insertions", alone or combined with "deletions". The deletions are produced when all of the k-tuples (or their majority) between two possible starts of the subfragments are false negatives. Since every position in the sequence is defined by k k-tuples, the occurrence of the deletions in a common case requires k consecutive false negatives. (With 10% of the false negatives and  $k=8$ , this situation takes place after every 108 elements). This situation is extremely infrequent even in mammalian genome sequencing using random libraries containing ten genome equivalents.

Elongation of the end of the sequence caused by false positive k-tuples is the special case of "insertions", since the end of the sequence can be considered as the endless linear array of false negative k-tuples. One may consider a group of false positive k-tuples producing subfragments longer than one k-tuple. Situations of this kind may be detected if subfragments are generated in overlapped fragments, like random physical fragments of the ordering library. An insertion, or insertion in place of a deletion, can arise as a result of specific combinations of false positive and false negative

WO 99/09217

PCT/US98/16966

k-tuples. In the first case, the number of consecutive false negatives is smaller than k. Both cases require several overlapping false positive k-tuples. The insertions and deletions are mostly theoretical possibilities without sizable practical repercussions since the requirements in the number and specificity of false k-tuples are simply too high.

- 5 In every other situation of not meeting the theoretical requirement of the minimal number an the kind of the false positive and/or negatives, mistakes in the k-tuples content may produce only the lesser completeness of a generated sequence.

SBH, a sample nucleic acid is sequenced by exposing the sample to a support-bound probe of known sequence and a labeled probe or probes in solution.

- 10 Wherever the probes ligase is introduced into the mixture of probes and sample, such that, wherever a support has a bound probe and a labeled probe hybridized back to back along the sample, the two probes will be chemically linked by the action of the ligase. After washing, only chemically linked support-bound and labeled probes are detected by the presence of the labeled probe. By knowing the identity of the support-bound probe at  
15 a particular location in an array, and the identity of the labeled probe, a portion of the sequence of the sample may be determined by the presence of a label at a point in an array on a Format with a sample of three substrate. And not chances not working are maximally overlapping sequences of all of the ligated probe pairs, the sequence of the sample may be reconstructed. Not of the sample to be sequenced may be a nucleic acid  
20 fragment or oligonucleotide of ten base pairs ("bp"). The sample is preferably four to one thousand bases in length.

- The length of the probe is a fragment less than ten bases in length, and, preferably, is between four and nine bases in length. In this way, arrays of support-bound probes may include all oligonucleotides of a given length or may include  
25 only oligonucleotides selected for a particular test. Where all oligonucleotides of a given length are used, the number of central oligonucleotides may be calculated by  $4^N$  where N is the length of the probe.

WO 99/09217

PCT/US98/16966

**EXAMPLE 18****Re-Using Sequencing Chips**

When ligation is employed in the sequencing process, then the ordinary oligonucleotide chip cannot be immediately reused. The inventor contemplates that this  
5 may be overcome in various ways.

One may employ ribonucleotides for the second probe, probe P, so that this probe may subsequently be removed by RNase treatment. RNase treatment may utilize RNase A an endoribonuclease that specifically attacks single-stranded RNA 3' to pyrimidine residues and cleaves the phosphate linkage to the adjacent nucleotide. The end products  
10 are pyrimidine 3' phosphates and oligonucleotides with terminal pyrimidine 3' phosphates. RNase A works in the absence of cofactors and divalent cations.

To utilize an RNase, one would generally incubate the chip in any appropriate RNase-containing buffer, as described by Sambrook *et al.* (1989; incorporated herein by reference). The use of 30-50 ul of RNase-containing buffer per 8 x 8 mm or 9 x 9 mm  
15 array at 37°C for between 10 and 60 minutes is appropriate. One would then wash with hybridization buffer.

Although not widely applicable, one could also use the uracil base, as described by Craig *et al.* (1989), incorporated herein by reference, in specific embodiments. Destruction of the ligated probe combination, to yield a re-usable chip, would be achieved  
20 by digestion with *the E. Coli* repair enzyme, uracil-DNA glycosylase which removes uracil from DNA.

One could also generate a specifically cleavable bond between the probes and then cleave the bond after detection. For example, this may be achieved by chemical ligation as described by Shabarova *et al.*, (1991) and Dolinnaya *et al.*, (1988), both references  
25 being specifically incorporated herein by reference.

Shabarova *et al.* (1991) describe the condensation of oligodeoxyribonucleotides with cyanogen bromide as a condensing agent. In their one step chemical ligation reaction, the oligonucleotides are heated to 97°C, slowly cooled to 0°C, then 1 ul 10 mM BrCN in acetonitrile is added.

30 Dolinnaya *et al.* (1988) show how to incorporate phosphoramidate and pyrophosphate internucleotide bonds in DNA duplexes. They also use a chemical ligation method for modification of the sugar phosphate backbone of DNA, with a water-soluble

WO 99/09217

PCT/US98/16966

carbodiimide (CDI) as a coupling agent. The selective cleavage of a phosphoamide bond involves contact with 15%  $\text{CH}_3\text{COOH}$  for 5 min at 95°C. The selective cleavage of a pyrophosphate bond involves contact with a pyridine-water mixture (9:1) and freshly distilled  $(\text{CF}_3\text{CO})_2\text{O}$ .

5

**EXAMPLE 19****Diagnostics - Scoring Known Mutations or Full Gene Resequencing**

In a simple case, the goal may be to discover whether selected, known mutations occur in a DNA segment. Less than 12 probes may suffice for this purpose, for example, 10 5 probes positive for one allele, 5 positive for the other, and 2 negative for both. Because of the small number of probes to be scored per sample, large numbers of samples may be analyzed in parallel. For example, with 12 probes in 3 hybridization cycles, 96 different genomic loci or gene segments from 64 patient may be analyzed on one 6 x 9 in membrane containing 12 x 24 subarrays each with 64 dots representing the same DNA 15 segment from 64 patients. In this example, samples may be prepared in sixty-four 96-well plates. Each plate may represent one patient, and each well may represent one of the DNA segments to be analyzed. The samples from 64 plates may be spotted in four replicas as four quarters of the same membrane.

A set of 12 probes may be selected by single channel pipetting or by a single pin 20 transferring device (or by an array of individually-controlled pipets or pins) for each of the 96 segments, and the selected probes may be arrayed in twelve 96-well plates. Probes may be labelled, if they are not prelabelled, and then probes from four plates may be mixed with hybridization buffer and added to the subarrays preferentially by a 96-channel pipetting device. After one hybridization cycle it is possible to strip off previously-applied 25 probes by incubating the membrane at 37° to 55°C in the preferably undiluted hybridization or washing buffer.

The likelihood that probes positive for one allele are positive and probes positive for the other allele are negative may be used to determine which of the two alleles is present. In this redundant scoring scheme, some level (about 10%) of errors in 30 hybridization of each probe may be tolerated.

An incomplete set of probes may be used for scoring most of the alleles, especially if the smaller redundancy is sufficient, e.g. one or two probes which prove the

WO 99/09217

PCT/US98/16966

presence or absence in a sample of one of the two alleles. For example, with a set of four thousand 8-mers there is a 91% chance of finding at least one positive probe for one of the two alleles for a randomly selected locus. The incomplete set of probes may be optimized to reflect G+C content and other biases in the analyzed samples.

- 5 For full gene sequencing, genes may be amplified in an appropriate number of segments. For each segment, a set of probes (about one probe per 2-4 bases) may be selected and hybridized. These probes may identify whether there is a mutation anywhere in the analyzed segments. Segments (i.e., subarrays which contain these segments) where one or more mutated sites are detected may be hybridized with additional probes to find
- 10 the exact sequence at the mutated sites. If a DNA sample is tested by every second 6-mer, and a mutation is localized at the position that is surrounded by positively hybridized probes TGCAA and TATTCC and covered by three negative probes: CAAAAC, AAATA and ACTATT, the mutated nucleotides must be A and/or C occurring in the normal sequence at that position. They may be changed by a single base
- 15 mutation, or by a one or two nucleotide deletion and/or insertion between bases AA, AC or CT.

- One approach is to select a probe that extends the positively hybridized probe TGCAA for one nucleotide to the right, and which extends the probe TATTCC one nucleotide to the left. With these 8 probes (GCAAAA, GCAAA, GCAAAC, GCAAAG
- 20 and ATATT, TATT, CTATT, GTATT) two questionable nucleotides are determined.

- The most likely hypothesis about the mutation may be determined. For example, A is found to be mutated to G. There are two solutions satisfied by these results. Either replacement of A with G is the only change or there is in addition to that change an
- 25 insertion of some number of bases between newly determined G and the following C. If the result with bridging probes is negative these options may then be checked first by at least one bridging probe comprising the mutated position (AAGCTA) and with an additional 8 probes: CAAAG, CAAAGT, CAAAGC, CAAAGG and ACTATT, TCTATT, CCTATT, GCTATT, I There are many other ways to select mutation-solving
- 30 probes.

In the case of diploid, particular comparisons of scores for the test samples and homozygotic control may be performed to identify heterozygotes (see above). A few

WO 99/09217

PCT/US98/16966

consecutive probes are expected to have roughly twice smaller signals if the segment covered by these probes is mutated on one of the two chromosomes.

#### **EXAMPLE 20**

##### **5 Identification of Genes (Mutations) Responsible for Genetic Disorders and Other Traits**

Using universal sets of longer probes (8-mers or 9-mers) on immobilized arrays of samples, DNA fragments as long as 5-20 kb may be sequenced without subcloning.

Furthermore, the speed of sequencing readily may be about 10 million

- 10 bp/day/hybridization instrument. This performance allows for resequencing a large fraction of human genes or the human genome repeatedly from scientifically or medically interesting individuals. To resequence 50% of the human genes, about 100 million bp is checked. That may be done in a relatively short period of time at an affordable cost.

- This enormous resequencing capability may be used in several ways to identify  
15 mutations and/or genes that encode for disorders or any other traits. Basically, mRNAs (which may be converted into cDNAs) from particular tissues or genomic DNA of patients with particular disorders may be used as starting materials. From both sources of DNA, separate genes or genomic fragments of appropriate length may be prepared either by cloning procedures or by in vitro amplification procedures (for example by PCR). If  
20 cloning is used, the minimal set of clones to be analyzed may be selected from the libraries before sequencing. That may be done efficiently by hybridization of a small number of probes, especially if a small number of clones longer than 5 kb is to be sorted. Cloning may increase the amount of hybridization data about two times, but does not require tens of thousands of PCR primers.

- 25 In one variant of the procedure, gene or genomic fragments may be prepared by restriction cutting with enzymes like Hga I which cuts DNA in following way: GACGC(N5')/CTGCG(N10'). Protruding ends of five bases are different for different fragments. One enzyme produces appropriate fragments for a certain number of genes. By cutting cDNA or genomic DNA with several enzymes in separate reactions, every gene of  
30 interest may be excised appropriately. In one approach, the cut DNA is fractionated by size. DNA fragments prepared in this way (and optionally treated with Exonuclease III which individually removes nucleotides from the 3' end and increases length and

WO 99/09217

PCT/US98/16966

specificity of the ends) may be dispensed in the tubes or in multiwell plates. From a relatively small set of DNA adapters with a common portion and a variable protruding end of appropriate length, a pair of adapters may be selected for every gene fragment that needs to be amplified. These adapters are ligated and then PCR is performed by universal  
5 primers. From 1000 adapters, a million pairs may be generated, thus a million different fragments may be specifically amplified in the identical conditions with a universal pair of primers complementary to the common end of the adapters.

If a DNA difference is found to be repeated in several patients, and that sequence change is nonsense or can change function of the corresponding protein, then the mutated  
10 gene may be responsible for the disorder. By analyzing a significant number of individuals with particular traits, functional allelic variations of particular genes could be associated by specific traits.

This approach may be used to eliminate the need for very expensive genetic mapping on extensive pedigrees and has special value when there is no such genetic data  
15 or material.

#### **EXAMPLE 21**

##### **Scoring Single Nucleotide Polymorphisms in Genetic Mapping**

Techniques disclosed in this application are appropriate for an efficient  
20 identification of genomic fragments with single nucleotide polymorphisms (SNUPs). In 10 individuals by applying the described sequencing process on a large number of genomic fragments of known sequence that may be amplified by cloning or by in vitro amplification, a sufficient number of DNA segments with SNUPs may be identified. The polymorphic fragments are further used as SNUP markers. These markers are either  
25 mapped previously (for example they represent mapped STSs) or they may be mapped through the screening procedure described below.

SNUPs may be scored in every individual from relevant families or populations by amplifying markers and arraying them in the form of the array of subarrays. Subarrays contain the same marker amplified from the analyzed individuals. For each marker, as in  
30 the diagnostics of known mutations, a set of 6 or less probes positive for one allele and 6 or less probes positive for the other allele may be selected and scored. From the significant association of one or a group of the markers with the disorder, chromosomal



WO 99/09217

PCT/US98/16966

position of the responsible gene(s) may be determined. Because of the high throughput and low cost, thousands of markers may be scored for thousands of individuals. This amount of data allows localization of a gene at a resolution level of less than one million bp as well as localization of genes involved in polygenic diseases. Localized genes may  
5 be identified by sequencing particular regions from relevant normal and affected individuals to score a mutation(s).

PCR is preferred for amplification of markers from genomic DNA. Each of the markers require a specific pair of primers. The existing markers may be convertible or new markers may be defined which may be prepared by cutting genomic DNA by Hga I  
10 type restriction enzymes, and by ligation with a pair of adaptors.

SNUP markers can be amplified or spotted as pools to reduce the number of independent amplification reactions. In this case, more probes are scored per one sample. When 4 markers are pooled and spotted on 12 replica membranes, then 48 probes (12 per marker) may be scored in 4 cycles.

15

## **EXAMPLE 22**

### **Detection and Verification of Identity of DNA Fragments**

DNA fragments generated by restriction cutting, cloning or in vitro amplification (e.g. PCR) frequently may be identified in a experiment. Identification may be  
20 performed by verifying the presence of a DNA band of specific size on gel electrophoresis. Alternatively, a specific oligonucleotide may be prepared and used to verify a DNA sample in question by hybridization. The procedure developed here allows for more efficient identification of a large number of samples without preparing a specific oligonucleotide for each fragment. A set of positive and negative probes may be selected  
25 from the universal set for each fragment on the basis of the known sequences. Probes that are selected to be positive usually are able to form one or a few overlapping groups and negative probes are spread over the whole insert.

This technology may be used for identification of STSs in the process of their mapping on the YAC clones. Each of the STSs may be tested on about 100 YAC clones  
30 or pools of YAC clones. DNAs from these 100 reactions possibly are spotted in one subarray. Different STSs may represent consecutive subarrays. In several hybridization cycles, a signature may be generated for each of the DNA samples, which signature

WO 99/09217

PCT/US98/16966

proves or disproves existence of the particular STS in the given YAC clone with necessary confidence.

To reduce the number of independent PCR reactions or the number of independent samples for spotting, several STSs may be amplified simultaneously in a reaction or PCR samples may be mixed, respectively. In this case more probes have to be scored per one dot. The pooling of STSs is independent of pooling YACs and may be used on single YACs or pools of YACs. This scheme is especially attractive when several probes labelled with different colors are hybridized together.

In addition to confirmation of the existence of a DNA fragment in a sample, the amount of DNA may be estimated using intensities of the hybridization of several separate probes or one or more pools of probes. By comparing obtained intensities with intensities for control samples having a known amount of DNA, the quantity of DNA in all spotted samples is determined simultaneously. Because only a few probes are necessary for identification of a DNA fragment, and there are N possible probes that may be used for DNA N bases long, this application does not require a large set of probes to be sufficient for identification of any DNA segment. From one thousand 8-mers, on average about 30 full matching probes may be selected for a 1000 bp fragment.

### **EXAMPLE 23**

#### **20 Identification of Infectious Disease Organisms and Their Variants**

DNA-based tests for the detection of viral, bacterial, fungal and other parasitic organisms in patients are usually more reliable and less expensive than alternatives. The major advantage of DNA tests is to be able to identify specific strains and mutants, and eventually be able to apply more effective treatment. Two applications are described below.

The presence of 12 known antibiotic resistance genes in bacterial infections may be tested by amplifying these genes. The amplified products from 128 patients may be spotted in two subarrays and 24 subarrays for 12 genes may then be repeated four times on a 8 x 12 cm membrane. For each gene, 12 probes may be selected for positive and negative scoring. Hybridizations may be performed in 3 cycles. For these tests, a much smaller set of probes is most likely to be universal. For example, from a set of one thousand 8-mers, on average 30 probes are positive in 1000 bp fragments, and 10 positive

WO 99/09217

PCT/US98/16966

probes are usually sufficient for a highly reliable identification. As described in Example 9, several genes may be amplified and/or spotted together and the amount of the given DNA may be determined. The amount of amplified gene may be used as an indicator of the level of infection.

- 5 Another example involves possible sequencing of one gene or the whole genome of an HIV virus. Because of rapid diversification, the virus poses many difficulties for selection of an optimal therapy. DNA fragments may be amplified from isolated viruses from up to 64 patients and resequenced by the described procedure. On the basis of the obtained sequence the optimal therapy may be selected. If there is a mixture of two virus
- 10 types of which one has the basic sequence (similar to the case of heterozygotes), the mutant may be identified by quantitative comparisons of its hybridization scores with scores of other samples, especially control samples containing the basic virus type only. Scores twice as small may be obtained for three to four probes that cover the site mutated in one of the two virus types present in the sample (see above).

15

#### **EXAMPLE 24**

##### **Forensic and Parental Identification**

- Sequence polymorphisms make an individual genomic DNA unique. This permits analysis of blood or other body fluids or tissues from a crime scene and comparison with
- 20 samples from criminal suspects. A sufficient number of polymorphic sites are scored to produce a unique signature of a sample. SBH may easily score single nucleotide polymorphisms to produce such signatures.

- A set of DNA fragments (10-1000) may be amplified from samples and suspects. DNAs from samples and suspects representing one fragment are spotted in one or several
- 25 subarrays and each subarray may be replicated 4 times. In three cycles, 12 probes may determine the presence of allele A or B in each of the samples, including suspects, for each DNA locus. Matching the patterns of samples and suspects may lead to discovery of the suspect responsible for the crime.

- The same procedure may be applicable to prove or disprove the identity of parents
- 30 of a child. DNA may be prepared and polymorphic loci amplified from the child and adults; patterns of A or B alleles may be determined by hybridization for each. Comparisons of the obtained patterns, along with positive and negative controls, aide in

WO 99/09217

PCT/US98/16966

the determination of familial relationships. In this case, only a significant portion of the alleles need match with one parent for identification. Large numbers of scored loci allow for the avoidance of statistical errors in the procedure or of masking effects of de novo mutations.

5

#### **EXAMPLE 25**

##### **Assessing Genetic Diversity of Populations or Species and Biological Diversity of Ecological Niches**

Measuring the frequency of allelic variations on a significant number of loci (for  
10 example, several genes or entire mitochondrial DNA) permits development of different  
types of conclusions, such as conclusions regarding the impact of the environment on the  
genotypes, history and evolution of a population or its susceptibility to diseases or  
extinction, and others. These assessments may be performed by testing specific known  
alleles or by full resequencing of some loci to be able to define de novo mutations which  
15 may reveal fine variations or presence of mutagens in the environment.

Additionally, biodiversity in the microbial world may be surveyed by resequencing  
evolutionarily conserved DNA sequences, such as the genes for ribosomal RNAs or genes  
for highly conservative proteins. DNA may be prepared from the environment and  
particular genes amplified using primers corresponding to conservative sequences. DNA  
20 fragments may be cloned preferentially in a plasmid vector (or diluted to the level of one  
molecule per well in multiwell plates and then amplified in vitro). Clones prepared this  
way may be resequenced as described above. Two types of information are obtained.  
First of all, a catalogue of different species may be defined as well as the density of the  
individuals for each species. Another segment of information may be used to measure the  
25 influence of ecological factors or pollution on the ecosystem. It may reveal whether some  
species are eradicated or whether the abundance ratios among species is altered due to the  
pollution. The method also is applicable for sequencing DNAs from fossils.

#### **EXAMPLE 26**

##### **Detection or Quantification of Nucleic Acid Species**

DNA or RNA species may be detected and quantified by employing a probe pair  
including an unlabeled probe fixed to a substrate and a labeled probe in a solution. The

WO 99/09217

PCT/US98/16966

species may be detected and quantified by exposure to the unlabeled probe in the presence of the labeled probe and ligase. Specifically, the formation of an extended probe by ligation of the labeled and unlabeled probe on the sample nucleic acid backbone is indicative of the presence of the species to be detected. Thus, the presence of label at a  
5 specific point in the array on the substrate after removing unligated labeled probe indicates the presence of a sample species while the quantity of label indicates the expression level of the species.

Alternatively, one or more unlabeled probes may be arrayed on a substrate as first members of pairs with one or more labeled probes to be introduced in solution.

10 According to one method, multiplexing of the label on the array may be carried out by using dyes which fluoresce at distinguishable wavelengths. In this manner, a mixture of cDNAs applied to an array with pairs of labeled and unlabeled probes specific for species to be identified may be examined for the presence of and expression level of cDNA species. According to a preferred embodiment this approach may be carried out to  
15 sequence portions of cDNAs by selecting pairs of unlabeled and labeled probes pairs comprising sequences which overlap along the sequence of a cDNA to be detected.

Probes may be selected to detect the presence and quantity of particular pathogenic organisms genome by including in the composition selected probe pairs which appear in combination only in target pathogenic genome organisms. Thus, while no single probe  
20 pair may necessarily be specific for the pathogenic organism genome, the combination of pairs is. Similarly, in detecting or sequencing cDNAs, it might occur that a particular probe is not be specific for a cDNA or other type of species. Nevertheless, the presence and quantity of a particular species may be determined by a result wherein a combination of selected probes situated at distinct array locations is indicative of the presence of a  
25 particular species.

An infectious agent with about 10kb or more of DNA may be detected using a support-bound detection chip without the use of polymerase chain reaction (PCR) or other target amplification procedures. According to other methods, the genomes of infectious agents including bacteria and viruses are assayed by amplification of a single target  
30 nucleotide sequence through PCR and detection of the presence of target by hybridization of a labelled probe specific for the target sequence. Because such an assay is specific for

WO 99/09217

PCT/US98/16966

only a single target sequence it therefore is necessary to amplify the gene by methods such as PCR to provide sufficient target to provide a detectable signal.

According to this example, an improved method of detecting nucleotide sequences characteristic of infectious agents through a Format 3-type reaction is provided wherein a solid phase detection chip is prepared which comprises an array of multiple different immobilized oligonucleotide probes specific for the infectious agent of interest. A single dot comprising a mixture of many unlabeled probes complementary to the target nucleic acid concentrates the label specific to a species at one location thereby improving sensitivity over diffuse or single probe labeling. Such multiple probes may be of overlapping sequences of the target nucleotide sequence but may also be non-overlapping sequences as well as non-adjacent. Such probes preferably have a length of about 5 to 12 nucleotides.

A nucleic acid sample exposed to the probe array and target sequences present in the sample will hybridize with the multiple immobilized probes. A pool of multiple labeled probes selected to specifically bind to the target sequences adjacent to the immobilized probes is then applied with the sample to an array of unlabeled oligonucleotide probe mixtures. Ligase enzyme is then applied to the chip to ligate the adjacent probes on the sample. The detection chip is then washed to remove unhybridized and unligated probe and sample nucleic acids and the presence of sample nucleic acid may be determined by the presence or absence of label. This method provides reliable sample detection with about a 1000-fold reduction of molarity of the sample agent.

As a further aspect of the invention, the signal of the labelled probes may be amplified by means such as providing a common tail to the free probe which itself comprises multiple chromogenic, enzymatic or radioactive labels or which is itself susceptible to specific binding by a further probe agent which is multiply labelled. In this way, a second round of signal amplification may be carried out. Labeled or unlabeled probes may be used in a second round of amplification. In this second round of amplification, a lengthy DNA sample with multiple labels may result in an increased amplification intensity signal between 10 to 100 fold which may result in a total signal amplification of 100,000 fold. Through the use of both aspects of this example, an

WO 99/09217

PCT/US98/16966

intensity signal approximately 100,000 fold may give a positive result of probe-DNA ligation without having to employ PCR or other amplification procedures.

According to a further aspect of the invention an array or super array may be prepared which consists of a complete set of probes, for example 4096 6-mer probes.

- 5 Arrays of this type are universal in a sense that they can be used for detection or partial to complete sequencing of any nucleic acid species. Individual spots in an array may contain single probe species or mixtures of probes, for example N(1-3) B(4-6) N(1-3) type of mixtures that are synthesized in the single reaction (N represents all four nucleotides, B one specific nucleotide and where the associated numbers are a range of
- 10 numbers of bases *i.e.*, 1-3 means "from one to three bases".) These mixtures provide stronger signal for a nucleic acid species present at low concentration by collecting signal from different parts of the same long nucleic acid species molecule. The universal set of probes may be subdivided in many subsets which are spotted as unit arrays separated by barriers that prevent spreading of hybridization buffer with sample and labeled probe(s).
- 15 For detection of a nucleic acid species with a known sequence one of more oligonucleotide sequences comprising both unlabelled fixed and labeled probes in solution may be selected. Labeled probes are synthesized or selected from the presynthesized complete sets of, for example, 7-mers. The labeled probes are added to corresponding unit arrays of fixed probes such that a pair of fixed and labeled probes will adjacently
- 20 hybridize to the target sequence such that upon administration of ligase the probes will be covalently bound.

- If a unit array contains more than one fixed probe (as separated spots or within the same spot) that are positive in a given nucleic acid species all corresponding labeled probes may be mixed and added to the same unit array. The mixtures of labeled probes
- 25 are even more important when mixtures of nucleic acid species are tested. One example of a complex mixture of nucleic acid species are mRNAs in one cell or tissue.

- According to one embodiment of the invention unit arrays of fixed probes allow use of every possible immobilized probe with cocktails of a relatively small number of labeled probes. More complex cocktails of labeled probes may be used if a multiplex
- 30 labeling scheme is implemented. Preferred multiplexing methods may use different fluorescent dyes or molecular tags that may be separated by mass spectroscopy.

WO 99/09217

PCT/US98/16966

Alternatively, according to a preferred embodiment of the invention, relatively short fixed probes may be selected which frequently hybridize to many nucleic acid sequences. Such short probes are used in combination with a cocktail of labeled probes which may be prepared such that at least one labeled probe corresponds to each of the  
5 fixed probes. Preferred cocktails are those in which none of the labeled probes corresponds to more than one fixed probe.

### **EXAMPLE 27**

#### **Interrogation of Segments of the HIV Virus with All Possible 10-mers**

10 In this example of Format III SBH, an array was generated on nylon membranes (e.g., Gene Screen) of all possible bound 5-mers (1024 possible pentamers). The bound 5-mer oligonucleotides were synthesized with 5' tails of 5'-TTTTTT-NNN-3' (N = all four bases A, C, G, T, at this step in the synthesis equal molar amounts of all four bases are added). These oligonucleotides were precisely spotted onto the nylon membrane, the  
15 spots were allowed to dry, and the oligonucleotides were immobilized by treating the dried spots with UV light. Oligonucleotide densities of up to 18 oligonucleotides per square nanometer were obtained using this method. After the UV treatment, the nylon membranes were treated with a detergent containing buffer at 60-80°C. The spots of oligonucleotides were gridded in subarrays of 10 by 10 spots, and each subarray has 64 5-  
20 mer spots and 36 control spots. 16 subarrays give 1024 5-mers which encompasses all possible 5-mers.

The subarrays in the array were partitioned from each other by physical barriers, e.g., a hydrophobic strip, that allowed each subarray to be hybridized to a sample without cross-contamination from adjacent subarrays. In a preferred embodiment, the  
25 hydrophobic strip is made from a solution of silicone (e.g., household silicone glue and seal paste) in an appropriate solvent (such solvents are well known in the art). This solution of silicone grease is applied between the subarrays to form lines which after the solvent evaporates act as hydrophobic strips separating the cells.

In this Format III example, the free or solution (nonbound) 5-mers were  
30 synthesized with 3' tails of 5'-NN-3' (N = all four bases A, C, G, T). In this embodiment, the free 5-mers and the bound 5-mers are combined to produce all possible 10-mers for sequencing a known DNA sequence of less than 20 kb. 20 kb of double



WO 99/09217

PCT/US98/16966

stranded DNA is denatured into 40 kb of single-stranded DNA. This 40 kb of ss DNA hybridizes to about 4% of all possible 10-mers. This low frequency of 10-mer binding and the known target sequence allow the pooling of free or solution (nonbound) 5-mers for treatment of each subarray, without a loss of sequence information. In a preferred  
5 embodiment, 16 probes are pooled for each subarray, and all possible 5-mers are represented in 64 total pools of free 5-mers. Thus, all possible 10-mers may be probed against a DNA sample using 1024 subarrays (16 subarrays for each pool of free 5-mers).

The target DNA in this embodiment represents two-600 bp segments of the HIV virus. These 600 bp segments are represented by pools of 60 overlapping 30-mers (the  
10 30-mers overlap each adjacent 30 mer by 20 nucleotides). The pools of 30-mers mimic a target DNA that has been treated using techniques well known in the art to shear, digest, and/or random PCR the target DNA to produce a random pool of very small fragments.

As described above in the previous Format III examples, the free 5-mers are labeled with radioactive isotopes, biotin, fluorescent dyes, etc. The labeled free 5-mers  
15 are then hybridized along with the bound 5-mers to the target DNA, and ligated. In a preferred embodiment, 300-1000 units of ligase are added to the reaction. The hybridization conditions were worked out following the teachings of the previous examples. Following ligation and removal of the target DNA and excess free probe, the array is assayed to determine the location of labeled probes (using the techniques  
20 described in the examples above).

The known DNA sequence of the target, and the known free and bound 5-mers in each subarray, predict which bound 5-mers will be ligated to a labeled free 5-mer in each subarray. The signal from 20 of these predicted dots were lost and 20 new signals were gained for each change in the target DNA from the predicted sequence. The overlapping  
25 sequence of the bound 5-mers in these ten new dots identifies which free, labeled 5-mer is bound in each new dot.

Using the described methods, arrays and pools of free, labeled 5-mers, the test HIV DNA sequence was probed with all possible 10-mers. Using this Format III approach, we properly identified the "wild-type" sequence of the segments tested, as well  
30 as several sequence "mutants" that were introduced into these segments.

WO 99/09217

PCT/US98/16966

**EXAMPLE 28****Sequencing of Repetitive DNA Sequences**

In one embodiment, repetitive DNA sequences in the target DNA are sequenced with "spacer oligonucleotides" in a modified Format III approach. Spacer

5 oligonucleotides of varying lengths of the repetitive DNA sequence (the repeating sequence is identified on a first SBH run) are hybridized to the target DNA along with a first known adjoining oligonucleotide and a second known, or group of possible oligonucleotides adjoining the other side of the spacer (known from the first SBH run). When a spacer matching the length of the repetitive DNA segment is hybridized to the

10 target, the two adjacent oligonucleotides can be ligated to the spacer. If the first known oligonucleotide is fixed to a substrate, and the second known or possible oligonucleotide(s) is labeled, a bound ligation product including the labeled second known or possible oligonucleotide(s) is formed when a spacer of the proper length is hybridized to the target DNA.

15

**EXAMPLE 29****Sequencing Through Branch Points with Format III SBH**

In one embodiment, branch points in the target DNA are sequenced using a third set of oligonucleotides and a modified Format III approach. After a first SBH run,

20 several branch points may be identified when the sequence is compiled. These can be solved by hybridizing oligonucleotide(s) that overlap partially with one of the known sequences leading into the branch point and then hybridizing to the target an additional oligonucleotide that is labeled and corresponds to one of the sequences that comes out of the branch point. When the proper oligonucleotides are hybridized to the target DNA,

25 the labeled oligonucleotide can be ligated to the other(s). In a preferred embodiment, a first oligonucleotide that is offset by one to several nucleotides from the branch point is selected (so that it reads into one of the branch sequences), a second oligonucleotide reading from the first and into the branch point sequence is also selected, and a set of third oligonucleotides that correspond to all the possible branch sequences with an overlap

30 of the branch point sequence by one or a few nucleotides (corresponding to the first oligonucleotide) is selected. These oligonucleotides are hybridized to the target DNA, and only the third oligonucleotide with the proper branch sequence (that matches the

WO 99/09217

PCT/US98/16966

branch sequence of the first oligonucleotide) will produce a ligation product with the first and second oligonucleotides.

### **EXAMPLE 30**

#### **5 Multiplexing Probes for Analyzing a Target Nucleic Acid**

In this Example, sets of probes are labeled with different labels so that each probe of a set can be differentiated from the other probes in the set. Thus, the set of probes may be contacted with target nucleic acid in a single hybridization reaction without the loss of any probe information. In preferred embodiments, the different labels are  
10 different radioisotopes, or different fluorescent labels, or different EMLs. These sets of probes may be used in either Format I, Format II or Format III SBH.

In Format I SBH, the set of differently labeled probes are hybridized to target nucleic acid which is fixed to a substrate under conditions that allow differentiation between perfect matches one base-pair mismatches. Specific probes which bind to the  
15 target nucleic acid are identified by their different labels and perfect matches are determined, at least in part, from this binding information.

In Format II SBH, the target nucleic acids are labeled with different probes and hybridized to arrays of probes. Specific target nucleic acids which bind to the probes are identified by their different labels and perfect matches are determined, at least in part,  
20 form this binding information.

In Format III SBH, the set of differently labeled probes and fixed probes are hybridized to a target nucleic acid under conditions that allow perfect matches to be differentiated from one base-pair mismatches. Labeled probes that are adjacent, on the target, to a fixed probe are bound to the fixed probe, and these products are detected and  
25 differentiated by their different labels.

In a preferred embodiment, the different labels are EMLs, which can be detected by electron capture mass spectrometry (EC-MS). EMLs may be prepared from a variety of backbone molecules, with certain aromatic backbones being particularly preferred, e.g., see Xu *et al.*, J. Chromatog. 764:95-102 (1997). The EML is attached to a probe  
30 in a reversible and stable manner, and after the probe is hybridized to target nucleic acid, the EML is removed from the probe and identified by standard EC-MS (e.g., the EC-MS may be done by a gas chromatograph-mass spectrometer).

WO 99/09217

PCT/US98/16966

**EXAMPLE 31****Detection of Low Frequency Target Nucleic Acids**

Format III SBH has sufficient discrimination power to identify a sequence that is present in a sample at 1 part to 99 parts of a similar sequence that differs by a single  
5 nucleotide. Thus, Format III can be used to identify a nucleic acid present at a very low concentration in a sample of nucleic acids, e.g., a sample derived from blood.

In one embodiment, the two sequences are for cystic fibrosis and the sequences differ from each other by a deletion of three nucleotides. Probes for the two sequences were as follows, probes distinguishing the deletion from wild type were fixed to a  
10 substrate, and a labeled contiguous probe was common to both. Using these targets and probes, the deletion mutant could be detected with Format III SBH when it was present at one part to ninety nine parts of the wild-type.

**EXAMPLE 32****15 Polaroid Apparatus and Method for Analyzing a Target Nucleic Acid**

An apparatus for analyzing a nucleic acid can be constructed with two arrays of nucleic acids, and an optional material that prevents the nucleic acids of the two arrays from mixing until such mixing is desired. The arrays of the apparatus may be supported by a variety of substrates, including but not limited to, nylon membranes, nitrocellulose  
20 membranes, or other materials disclosed above. In preferred embodiments, one of the substrate is a membrane separated into sectors by hydrophobic strips, or a suitable support material with wells which may contain a gel or sponge. In this embodiment, probes are placed on a sector of the membrane, or in the well, the gel, or sponge, and a solution (with or without target nucleic acids) is added to the membrane or well so that  
25 the probes are solubilized. The solution with the solubilized probes is then allowed to contact the second array of nucleic acids. The nucleic acids may be, but are not limited to, oligonucleotide probes, or target nucleic acids, and the probes or target nucleic acids may be labeled. The nucleic acids may be labeled with any labels conventionally used in the art, including but not limited to radioisotopes, fluorescent labels or electrophore mass  
30 labels.

The material which prevents mixing of the nucleic acids may be disposed between the two arrays in such a way that when the material is removed the nucleic acids of the

WO 99/09217

PCT/US98/16966

two arrays mix together. This material may be in the form of a sheet, membrane, or other barrier, and this material may be comprised of any material that prevents the mixing of the nucleic acids.

This apparatus may be used in Format I SBH as follows: a first array of the  
5 apparatus has target nucleic acids that are fixed to the substrate, and a second array of the apparatus has nucleic acid probes that are labeled and can be removed to interrogate the target nucleic acid of the first array. The two arrays are optionally separated by a sheet of material that prevents the probes from contacting the target nucleic acid, and when this sheet is removed the probes can interrogate the target. After appropriate incubation and  
10 (optionally) washing steps the array of targets may be "read" to determine which probes formed perfect matches with the target. This reading may be automated or can be done manually (e.g., by eye with an autoradiogram). In Format II SBH, the procedure followed would be similar to that described above except that the target is labeled and the probes are fixed.

15 Alternatively, the apparatus may be used in Format III SBH as follows: two arrays of nucleic acid probes are formed, the nucleic acid probes of either or both arrays may be labeled, and one of the arrays may be fixed to its substrate. The two arrays are separated by a sheet of material that prevents the probes from mixing. A Format II reaction is initiated by adding target nucleic acid and removing the sheet allowing the probes to mix  
20 with each other and the target. Probes which bind to adjacent sites on the target are bound together (e.g., by base-stacking interactions or by covalently joining the backbones), and the results are read to determine which probes bound to the target at adjacent sites. When one set of probes is fixed to the substrate, the fixed array can be read to determine which probes from the other array are bound together with the fixed  
25 probes. As with the above method, this reading may be automated (e.g., with an ELISA reader) or can be done manually (e.g., by eye with an autoradiogram).

### **EXAMPLE 33**

#### **Three Dimensional Arrays Of Probes**

30 In a preferred embodiment, the oligonucleotide probes are fixed in a three-dimensional array. The three-dimensional array is comprised of multiple layers, such that each layer may be analyzed separate and apart from the other layers, or all the layers of

WO 99/09217

PCT/US98/16966

the three-dimensional array may be simultaneously analyzed. Three dimensional arrays include, for example, an array disposed on a substrate having multiple depressions with probes located at different depths within the depressions (each level is made up of probes at similar depths within the depression); or an array disposed on a substrate having  
5 depressions of different depths with the probes located at the bottom of the depression, at the peaks separating the depressions or some combination of peaks and depressions (each level is made up of all probes at a certain depth); or an array disposed on a substrate comprised of multiple sheets that are layered to form a three-dimensional array.

Materials for synthesizing these three-dimensional arrays are well known in the  
10 art, and include the materials previously recited in this specification as suitable as supports for probe arrays. In addition, other suitable materials which can support oligonucleotide probes, and which preferably, are flexible may be used as substrates.

#### **EXAMPLE 34**

##### **15 Signature Processing For Clustering cDNA Clones**

A plurality of distinct nucleic acid sequences were obtained from cDNA library, using standard pcr, SBH sequence signature analysis and Sanger sequencing techniques. The inserts of the library were amplified with pcr using primers specific for vector sequences which flank the inserts. These samples were spotted onto nylon membranes  
20 and interrogated with suitable number of oligonucleotide probes and the intensity of positive binding probes was measured giving sequence signatures. The clones were clustered into groups of similar or identical sequence signatures, and single representative clones were selected from each group for gel sequencing. The 5' sequence of the amplified inserts was then deduced using the reverse M13 sequencing primer in a typical  
25 Sanger sequencing protocol. PCR products were purified and subjected to fluorescent dye terminator cycle sequencing. Single pass gel sequencing was done using a 377 Applied Biosystems (ABI) sequencer. The majority of clones which were selected and sequenced by this method had sequences which differed from each other, and a very small number had the same sequence.

30

WO 99/09217

PCT/US98/16966

**EXAMPLE 35****High-Throughput Production Of Chips**

In a preferred embodiment, an apparatus for mass producing arrays of probes may comprise a rotating drum or plate coupled with an ink-jet deposition apparatus, for example, a microdrop dosing head; and a suitable robotics systems, for example, an anorad gantry. A particularly preferred embodiment of the apparatus will be described referring to Figs. 1-3.

The apparatus comprises a cylinder (1) to which a suitable substrate is fixed. The substrate may be any of the materials previously described as suitable for an array of probes. In a preferred embodiment, the substrate is a flexible material, and the arrays are made directly on the substrate. In alternative embodiments, a flexible substrate is fixed to the cylinder and individual chips are fixed on the substrate. The arrays are then made on each individual chip.

In a preferred embodiment, physical barriers are applied to the substrate or chip and define an array of wells. The physical barriers may be applied to the substrate or chip by the apparatus, or alternatively, the physical barriers are applied to the chips or substrate before they are fixed to the cylinder (1). A single spot of oligonucleotide probes is then placed into each well, wherein the probes placed into an individual well may all have the same sequence, or the probes spotted into an individual well may have different sequences. In a more preferred embodiment, the probe or probes spotted into each individual well in an array are different from the probe or probes spotted in the other wells of the array. Sequencing chips comprising multiple arrays can then be assembled from these arrays.

After the substrate or substrate and chips are fixed to the cylinder (1), a motor (not shown) rotates the cylinder. The cylinder's rotation speed is precisely determined by any of the ways well known in the art, including, for example using a fixed optical sensor and light source that rotates with the cylinder. A dispensing apparatus (3) moves along an arm (2) and can deliver probes or other reagents through a dispensing tip (8) to precise locations on the substrate or chips using the precise rotation speed calculated above, by methods well known in the art. The dispensing apparatus receives probes or reagents from the reservoir (6) through the feeding line (7). The reservoir (6) holds all the necessary probes and other reagents for making the arrays.

WO 99/09217

PCT/US98/16966

The dispensing apparatus is depicted in Figure 3. The dispensing apparatus may have one or multiple dispensing tips (14 & 8). Each dispensing tip has a sample well (13) in a body (12) that receives probes or other reagents through a sample line (10). The pressure line (11) pressurizes the chamber (9) to a psi sufficient to force probes or reagents through the dispensing tip (14 & 8). The sample line (10), well (13) and dispensing tip (14 & 8) must be flushed between each change in probe or reagent. An appropriate washing buffer is supplied through sample line (10) or through an optional dedicated washing line (not shown) to the sample well (13) or optionally a portion or all of the chamber (9) may be filled with washing buffer. The washing buffer is then removed from the sample well (13) and chamber (9) if necessary by an evacuation line (not shown) or through the sample line (10) and dispensing tip (14 & 8).

When the dispensing means has applied probes to all the appropriate sites in each array or chip, the substrate (with or without chips) is removed from the cylinder and a new substrate is fixed to the cylinder.

#### **EXAMPLE 36**

##### **Analysis Of A Target Nucleic Acid With Probes Complexed To Discrete Particles**

In this embodiment, a target nucleic acid is interrogated with probes that are complexed (covalent or noncovalent) to a plurality of discrete particles. The discrete particles can be discriminated from each other based on a physical property (or a combination of physical properties), and particles with differentiated by the physical property are complexed with different probes. In a preferred embodiment, the probe is an oligonucleotide of a known sequence and length. Thus, a probe may be identified by the physical property of the discrete particle. Suitable probes for this embodiment include all the probes that are described above in previous sections, including probes which are shorter in an informative sense than the probes full length.

The physical property of the discrete particle may be any property, well known in the art, which allows particles to be differentiated into sets. For example, the particles could be differentiated into sets based on their size, fluorescence, absorbance, electromagnetic charge, or weight, or the particles could be labeled with dyes, radionuclides, or EMLs. Other suitable labels include ligands which can serve as specific binding members to a labeled antibody, chemilumescers, enzymes, antibodies which can



WO 99/09217

PCT/US98/16966

serve as a specific binding pair member for a labeled ligand, and the like. A wide variety of labels have been employed in immunoassays which can readily be employed. Still other labels include antigens, groups with specific reactivity, and electrochemically detectable moieties. Still further labels, include any of the labels recited above in  
5 previous sections. These labels and properties may be measured quantitatively by methods well known in the art, including for example, those methods described above in previous sections, and the particles may be differentiated on the basis of signal intensity or signal type (for one of the labels, e.g., different dye densities may be applied to a particle, or different types of dyes). In a preferred embodiment, several physical  
10 properties are combined and the different combinations of properties allow discrimination of the particles (e.g., ten sizes and ten colors could be combined to differentiate 100 particle groups).

The particle-probes allow the exploitation of standard combinatorial approaches so that, for example, all possible 10-mers can be synthesized using about 2000 reaction  
15 containers. A first set of 1024 reactions are done to synthesize all possible 5-mers on 1024 differentially labeled particles. The resulting probe-particles are mixed together, and split into another set of 1024 reaction containers. A second set of reactions are done with these samples to synthesize all possible 5-mer extensions on the probes in the pools of particles. The physical property identifies the first five nucleotides of each probe and  
20 the reaction container will identify the identity of the second five nucleotides of every probe. Thus, all possible 10-mer probes are synthesized using 2048 reaction containers. This approach is easily modified to make all possible n-mers for a large range of probe lengths.

In a preferred embodiment, the particles are separated into sets by the intensity of  
25 fluorescence of the particles. The particles in each set are prepared with varying densities of fluorescent label, and thus, the particles have different fluorescence intensities. The fluorescent intensity of fluorescein is related to concentration over a range of 1:300 to 1:300,000 (Lockhart et al., 1986), and between 1:3000 to 1:300,000 there is a linear relationship (so the fluorescein intensity is linear over a range of about 1-300). In the  
30 linear range of detection, 256 sets of particles are labeled with fluorescein (e.g., 3-259). 256 sets of particles allows all possible 4-mers to be attached to different sets of particles. By pooling the particles, all possible 5-mers can be made by having four pools of all

WO 99/09217

PCT/US98/16966

possible 4-mers and then extending the probes in each pool by A, G, C, or T. Similarly, all possible 6-mers can be made by having 16 pools of all possible 4-mers in which each pool of 4-mers is extended by one of the 16 possible two base permutations of A, G, C, and T (etc. for 7-mers there are 64 pools, 8-mers there are 256 pools, and so forth).

- 5        The 5-mer probes (in four pools) are used to interrogate a target nucleic acid. The target nucleic acid is labeled with another fluorescent dye, or other different label (as described above). Labeled target is mixed with the four pools, and complementary probes in each pool hybridize to the target nucleic acid. These hybridization complexes are detected by methods well-known in the art, and the positively hybridizing probes are
- 10 then identified by detecting the fluorescence intensity of the particle. In a most preferred embodiment, the mixture of probe-particles and target nucleic acid are fed through a flow-cytometer or other separating instrument one particle at a time, and the particle label and the target are measured to determine which probes are complementary to the target nucleic acid.
- 15        In an alternative embodiment, a set of free probes is labeled with another fluorescent dye, or other label (as described above), and individual free probes are mixed with each pool of 5-mer probes (four pools) and then the mixtures are hybridized with the target nucleic acid. An agent is added to covalently attach free probe to 5-mer probe (see previous sections for a description of suitable agents), when the free probe is bound to a
- 20 site on the target nucleic acid that is adjacent to the site on which the 5-mer probe is bound (the free probe site must be adjacent the end of the 5-mer probe which can be ligated). The particles are then assayed, by methods well known in the art, to determine which particles have been covalently coupled to the free probe, i.e., the particles which have the free probe label, and the 5-mer probe is identified by the fluorescent intensity of
- 25 the particle. In a most preferred embodiment, the mixture of probe-particles, free probes, and target nucleic acid are fed through a flow-cytometer one particle at a time, and the particle label and the free probe label are measured to determine which probes are complementary to the target nucleic acid.

- In preferred embodiment, a single apparatus houses all or most of the
- 30 manipulations for an analysis of a target nucleic acid with the probe-particle complexes. The apparatus has one or more reagent chambers in which buffer and labeled target nucleic acid are thoroughly mixed (target nucleic acid may be added manually or

WO 99/09217

PCT/US98/16966

automatically). The mixture is aliquoted from the reagent chamber into a plurality of reaction chambers, and each reaction chamber has a pool of probe-particle complexes. The probe-particles and target nucleic acid react under conditions which allow complementary probes to bind with the target nucleic acid. Excess target nucleic acid, 5 i.e., nonbound, is removed from the reaction chamber (e.g., by washing), and the particles bound to target nucleic acid are identified by the association of target nucleic acid label with the particles, and the probe is identified by the physical property of the particle. In a preferred embodiment, after removing excess target, the particles move single file through a channel from the reaction chamber to the detecting device(s). As 10 single particles move past the detecting device(s) they measure the target label and the physical property of the particle. In an alternative preferred embodiment, before or after removing excess target, the particles are fractionated, for example, by size (e.g., exclusion chromatography), charge (e.g., ion exchange chromatography), and/or density-weight into their sets using one or a combinations of these physical properties. These 15 fractionated particles are then assayed by the detecting device(s).

In an alternative of this embodiment, the main reagent chamber is supplied with buffer, target nucleic acid, a pool of probe-particle complexes, and a chemical or enzymatic ligating reagent. These components are thoroughly mixed and then aliquoted from the reagent chamber into a plurality of reaction chambers. Each of the reaction 20 chambers has a labeled free probe. Alternatively, the pool of probe particle complexes may be placed in the reaction chamber with the free probe instead of adding them to the reagent chamber. Additionally, the free probes could be added to the reagent chamber, and the pool of probe particles could be added to the reaction chamber. The probe-particles, target nucleic acid, and free probe react under conditions which allow free- and 25 particle-probes to bind with adjacent sites on the target nucleic acid so that free probe is ligated to the probe particle. Excess free probe (i.e., nonligated), and target nucleic acid are removed from the reaction chamber (e.g., by washing), and the ligated probes are identified by the association of free probe label with the particles, and the probes complexed to the particles are identified by the physical property of the particle. In a 30 preferred embodiment, after removing excess probe and the target, the particles move single file through a channel from the reaction chamber to the detecting device(s). As single particles move past the detecting device(s) they measure free probe label covalently

WO 99/09217

PCT/US98/16966

attached to the particles, and the physical property of the particle. In an alternative preferred embodiment, before or after removing excess probe and the target, the particles are fractionated, for example, by size (e.g., exclusion chromatography), charge (e.g., ion exchange chromatography), and/or density/weight into their sets using the physical  
5 property. These fractionated particles are assayed by the detecting device(s).

In a preferred embodiment, there is a set of second reaction chambers in the apparatus, and the pool of probe-particles are placed in the second reaction chamber. Target and buffer are mixed in the reagent chamber and these are fed into the first reaction chamber which contains the labeled free probe. The probe and target are mixed,  
10 and optionally the probe may hybridize to the target. This mixture of labeled probe and target is then passed to the second reaction chamber which contains the pool of probe-particles. The free probe and probe-particles hybridize to target and appropriate probes are ligated in the second reaction chamber. The ligating agent may be added at the reagent chamber or in either reaction chamber, preferably, the ligating agent is added in  
15 the second reaction chamber. The probe-particle hybridization products in the second reaction chamber are analyzed as above.

In one embodiment, the target nucleic acid is not amplified prior to analysis (either by PCR or in a vector, e.g., a lambda library). Preferably longer free and particle probes are used in this embodiment because of the increase in sequence complexity of the  
20 sample (i.e., to distinguish positives over background).

The probe-particle embodiments described in this example are suitable for use in any of the applications previously described, including, but not limited to the previously described diagnostic and sequencing applications. Additionally, these probe particle embodiments may be modified by any the previously described variations or  
25 modifications.

### **EXAMPLE 37**

#### **The Interaction of Complementary Polynucleotides in the Presence of Agents Which Modify the Binding Between the Polynucleotides.**

30 In this embodiment, the discrimination of perfect matches from mismatches in the binding of complementary polynucleotides is modulated by the addition of an agent or agents. In a preferred embodiment, the complementary polynucleotides are a target

WO 99/09217

PCT/US98/16966

polynucleotide and a polynucleotide probe. The discrimination of perfect matches from mismatches may be modulated by adding an agent wherein the agent is a salt such as trialkyl ammonium salt (e.g., TMAC, Ricelli et al., Nucl. Acids Res. 21:3785-3788 (1993)), sodium chloride, phosphate salts, and borate salts, organic solvents such as  
5 formamide, glycol, dimethylsulfoxide, and dimethylformamide, urea, guanidinium, amino acid analogs such as betaine (Henke et al., Nucl. Acids Res. 19:3957-3958 (1997); Rees et al., Biochemistry 32:137-144 (1993)), polyamines such as spermidine and spermine (Thomas et al., Nucl. Acids Res. 25:2396-2402 (1997)), or other positively charged molecules which neutralize the negative charge of the phosphate backbone, detergents  
10 such as sodium dodecyl sulfate, and sodium lauryl sarcosinate, minor/major groove binding agents, positively charged polypeptides, and intercalating agents such as acridine, ethidium bromide, and anthracine. In a preferred embodiment, a mixture of agents is added to the hybridization reaction to modulate the discrimination of perfect matches from mismatches. Some of these agents effect discrimination by reducing the entropy of  
15 melting between two complementary strands.

In a preferred embodiment, the discrimination of perfect matches from mismatches is improved by the agent or agents. For example, formamide, a commonly used denaturing agent, has been shown to preferentially destabilize mismatches versus perfect matches in a format III reaction. As described above, a format III reaction was set up,  
20 and then varying amounts of formamide were added (0%, 10%, 20%, 30%, 40%, and 50%). At 0% a perfect match signal was detected and the background (mismatches) was high. At 10% formamide, there was a good perfect match signal and the background/mismatch signal was reduced. At 20% formamide, the perfect match signal was reduced (but detectable) and the background/mismatch signal was eliminated. At  
25 30%-50% formamide there was no perfect match or background/mismatch signal.

In an alternative embodiment, an agent is used to reduce or increase the  $T_m$  of a pair of complementary polynucleotides. In a more preferred embodiment, a mixture of the agents is used to reduce or increase the  $T_m$  of a pair of complementary polynucleotides. The agents may alter the  $T_m$  in a number of ways, two examples, which  
30 are not meant to limit the invention, are (1) agents which disrupt the hydrogen bonding between the bases of two complementary polynucleotides (Goodman, Proc. Nat'l Acad. Sci. 94:10493-10495 (1997); Moran et al., Proc. Nat'l Acad. Sci. 94:10506-10511

WO 99/09217

PCT/US98/16966

(1997); Nguyen et al., Nucl. Acids Res. 25:3059-3065 (1997)), and (2) agents which neutralize or shield the negative charges of the phosphates in the sugar phosphate backbone of the polynucleotide (Thomas et al., Nucl. Acids Res. 25:2396-2402 (1997)).

By strengthening or weakening (1) and/or (2) one can modulate the  $T_m$  of a

5 complementary pair of polynucleotides.

In a most preferred embodiment, an agent or agents are added to decrease the binding energy of GC base pairs, or increase the binding energy of AT base pairs, or both. In a preferred embodiment, the agent or agents are added so that the binding energy from an AT base pair is approximately equivalent to the binding energy of a GC

10 base pair. Thus, the energy of binding between two complementary polynucleotides is solely dependent on length. The energy of binding of these complementary polynucleotides may be increased by adding an agent that neutralizes or shields the negative charges of the phosphate groups in the polynucleotide backbone.

15 The present invention is not to be limited in scope by the exemplified embodiments which are intended as illustrations of single aspects of the invention, and compositions and methods which are functionally equivalent are within the scope of the invention. Indeed, numerous modifications and variations in the practice of the invention are expected to occur to those skilled in the art upon consideration of the present

20 preferred embodiments. Consequently, the only limitations which should be placed upon the scope of the invention are those which appear in the appended claims.

All references cited within the body of the instant specification are hereby incorporated by reference in their entirety.

WO 99/09217

PCT/US98/16966

CLAIMS

## WHAT IS CLAIMED IS:

1. An array of a plurality of oligonucleotide probes, comprising:  
5 a substrate;  
a material that forms a physical barrier, wherein the material is disposed on the substrate to form a grid having a plurality of wells; and  
wherein the plurality of oligonucleotide probes are arranged to form an array in the plurality of wells, wherein each well contains one spot of probes fixed on the  
10 substrate.
2. The array of Claim 1, wherein each individual spot has probes with sequences that are different from the other probes at the other spots in the array.
- 15 3. The array of Claim 2, wherein the probes in each individual spot have the same sequence.
4. The array of Claim 1, wherein the distance between a center of one spot and a center of an adjacent spot is at least 325  $\mu\text{m}$ .  
20
5. A sequencing chip comprising a plurality of arrays of Claim 1.
6. A sequencing chip comprising a plurality of arrays of Claim 2.
- 25 7. A sequencing chip comprising a plurality of arrays of Claim 3.
8. The array of Claim 1, wherein the oligonucleotide probe comprises an informational portion and a reactive group wherein the reactive group is used to attach the probe to the substrate.  
30
9. The array of Claim 8, wherein the oligonucleotide probe further comprises at least one randomized position.

WO 99/09217

PCT/US98/16966

10. The array of Claim 9, wherein the oligonucleotide probe further comprises a spacer.

11. An array of a plurality of oligonucleotide probes, comprising a substrate  
5 having a plurality of levels, wherein the plurality of probes are fixed to the plurality of levels in the substrate.

12. The array of Claim 11, wherein each level may be analyzed individually.

10 13. The array of Claim 11, wherein the plurality of levels may be analyzed simultaneously.

14. The array of Claim 11, wherein the oligonucleotide probe comprises an informational portion and a reactive group wherein the reactive group is used to attach the  
15 probe to the substrate.

15. The array of Claim 14, wherein the oligonucleotide probe further comprises at least one randomized position.

20 16. The array of Claim 15, wherein the oligonucleotide probe further comprises a spacer.

17. A method for analyzing a target nucleic acid, comprising the steps of:  
contacting the target nucleic acid with a plurality of oligonucleotide probes,  
25 wherein the probes are complexed with a plurality of different discrete particles that can be discriminated from one another based on a physical property and wherein a different probe is complexed with each type of discrete particle;  
detecting those probes that are complementary to the target nucleic acid; and  
analyzing the target nucleic acid from a set of complementary probes.

30



WO 99/09217

PCT/US98/16966

18. The method of Claim 17, further comprising the step of separating the discrete particles into fractions, wherein the discrete particles are separated on the basis of the physical property.

5 19. The method of Claim 18, wherein the set of complementary probes has at least two probes which overlap.

20. The method of Claim 18, wherein a sequence of the target nucleic acid is compiled in the analyzing step.

10

21. The method of Claim 18, wherein the complementary probes are identified by the physical property of the discrete particle.

22. The method of Claim 18, wherein wherein the physical property is  
15 associated with a molecule selected from the group consisting of a dye, a radionucleotide, an EML, and a flourescent molecule.

23. The method of Claim 18, wherein the physical property is selected from the group comprising a size, a charge, an absorbance, and a weight.

20

24. The method of Claim 23, wherein the physical property is associated with an intensity of the physical property.

25. The method of Claim 23, wherein the physical property is associated with a  
25 plurality of different molecules.

26. The method of Claim 18, wherein an informative portion of the probes is shorter than the probes full length.

30 27. The method of Claim 18, wherein the target nucleic acid contains a label, and the complementary probes are detected by the label on the target nucleic acid.

WO 99/09217

PCT/US98/16966

28. The method of Claim 17, wherein the detecting step is performed on individual discrete particles by passing the individual particles past a detector.

29. The method of Claim 28, wherein the set of complementary probes has at  
5 least two probes which overlap.

30. The method of Claim 28, wherein a sequence of the target nucleic acid is compiled in the analyzing step.

10 31. The method of Claim 28, wherein the complementary probes are identified by the physical property of the discrete particle.

32. The method of Claim 28, wherein wherein the physical property is associated with a molecule selected from the group consisting of a dye, a radionucleotide,  
15 an EML, and a flourescent molecule.

33. The method of Claim 28, wherein the physical property is selected from the group comprising a size, a charge, an absorbance, and a weight.

20 34. The method of Claim 33, wherein the physical property is associated with an intensity of the physical property.

35. The method of Claim 33, wherein the physical property is associated with a plurality of different molecules.

25

36. The method of Claim 28, wherein an informative portion of the probes is shorter than the probes full length.

37. The method of Claim 28, wherein the target nucleic acid contains a label,  
30 and the complementary probes are detected by the label on the target nucleic acid.

WO 99/09217

PCT/US98/16966

38. The method of Claim 28, further comprising the step of contacting the target nucleic acid with a plurality of free oligonucleotides, and

covalently joining a complementary free probe, bound at a site in the target nucleic acid, to a complementary probe complexed to the discrete particle which is bound to a  
5 site on the target nucleic acid that is adjacent to the site on which the free probe is bound, and

wherein the detecting step identifies the free probe which is covalently joined to the probe of the discrete particle.

10 39. The method of Claim 38, wherein the set of complementary, covalently joined probes has at least two covalently joined probes which overlap.

40. The method of Claim 38, wherein a sequence of the target nucleic acid is compiled in the analyzing step.

15

41. The method of Claim 38, wherein the probes complexed to the discrete particle are identified by the physical property of the discrete particle.

42. The method of Claim 38, further comprising the step of separating the  
20 discrete particles into fractions, wherein the discrete particles are separated on the basis of the physical property.

43. The method of Claim 42, wherein the discrete particles are separated into fractions using a flow cytometer.

25

44. The method of Claim 38, wherein the physical property is associated with a molecule selected from the group consisting of a dye, a radionucleotide, an EML, and a fluorescent molecule.

WO 99/09217

PCT/US98/16966

45. The method of Claim 38, wherein the physical property property is selected from the group comprising a size, a charge, an absorbance, and a weight.

46. The method of Claim 45, wherein the physical property is associated with  
5 an intensity of the physical property.

47. The method of Claim 45, wherein the physical property is associated with a plurality of different molecules.

10 48. The method of Claim 38, wherein an informative portion of the free probes is shorter than the probes full length.

49. The method of Claim 38, wherein an informative portion of the probes complexed to the discrete particles is shorter than the probes full length.

15

50. The method of Claim 38, wherein an informative portion of the free probes and an informative portion of the probes complexed to the discrete particles are shorter than the probes full length.

20 51. A method for analyzing a target nucleic acid, comprising the steps of:  
contacting the target nucleic acid with a probe under conditions which allow a perfect match to be discriminated from a mismatch, wherein an agent is added which increase the discrimination of the perfect match from the mismatch; and  
detecting whether the probe is complementary to the target nucleic acid.

25

52. The method of Claim 51, wherein the agent is selected from the group comprising a salt, an organic solvent, an urea, a guanidinium, an amino acid analogs, a

WO 99/09217

PCT/US98/16966

polyamine, other positively charged molecules which neutralize the negative charge of the phosphate backbone, a detergent, a minor/major groove binding agent, a positively charged polypeptide, and an intercalating agent.

5           53.    The method of Claim 52, wherein the salt is selected from the group comprising a trialkyl ammonium salt, a sodium chloride, a phosphate salt, and a borate salt.

          54.    The method of Claim 52, wherein the organic solvent is selected from the  
10 group comprising a formamide, a glycol, a dimethylsulfoxide, and a dimethylformamide.

          55.    The method of Claim 52, wherein the amino acid analog is a betaine.

          56.    The method of Claim 52, wherein the polyamine is selected from the group  
15 comprising a spermidine and a spermine.

          57.    The method of Claim 52, wherein the detergent is selected from the group comprising a sodium dodecyl sulfate, and a sodium lauryl sarcosinate.

20           58.    The method of Claim 52, wherein the intercalating agent is selected from the group comprising an acridine, an ethidium bromide, and an anthracine.

          59.    The method of Claim 51, wherein a plurality of agents is added.

25           60.    The method of Claim 59, wherein the agents are selected from the group comprising a salt, an organic solvent, an urea, a guanidinium, an amino acid analogs, a polyamine, other positively charged molecules which neutralize the negative charge of the

WO 99/09217




PCT/US98/16966

phosphate backbone, a detergent, a minor/major groove binding agent, a positively charged polypeptide, and an intercalating agent.

61. A method for analyzing a target nucleic acid, comprising the steps of:
- 5 providing an array of a plurality of fixed oligonucleotide probes;
- providing a plurality of labeled oligonucleotide probes;
- contacting the target nucleic acid with the fixed probes and the labeled probes under conditions that allow the probes which form perfect matches with the target nucleic acid to be distinguished from the probes bound with a one base mismatch to the target
- 10 nucleic acid, wherein an agent is added which increase the discrimination of the perfect match from the one base pair mismatch;
- covalently joining the fixed probe, bound at a site in the target nucleic acid, to the labeled probe, which is hybridized to a site on the target nucleic acid that is adjacent to the site on which the fixed probe is bound; and
- 15 identifying the fixed probes and the labeled probes that are covalently joined.
62. The method of Claim 61, wherein the agent is selected from the group comprising a salt, an organic solvent, an urea, a guanidinium, an amino acid analogs, a polyamine, other positively charged molecules which neutralize the negative charge of the
- 20 phosphate backbone, a detergent, a minor/major groove binding agent, a positively charged polypeptide, and an intercalating agent.
63. The method of Claim 61, wherein the agent is formamide.
- 25 64. The method of Claim 61, wherein a plurality of agents is added.


WO 99/09217

PCT/US98/16966



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/16966

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(6) : C12Q 1/68; C12P 19/34; C07H 21/04 US CL : 435/6, 91.2; 536/24.3 According to International Patent Classification (IPC) or to both national classification and IPC														
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/6, 91.2; 536/24.3 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched NONE Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Please See Extra Sheet														
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>														
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.												
Y	WO 94/27719 A1 (BRENNAN, THOMAS, M.) 08 December 1994, pages 10 and 12.	1-16												
Y	US 5,545,531 A (RAVA et al) 13 August 1996, column 8.	1-16												
Y	WO 92/10588 A1 (AFFYMAX TECHNOLOGIES N.V.) 25 June 1992, pages 30-31	17-50												
Y	US 5,585,241 A (LINDMO) 17 December 1996, columns 4 and 8.	17-50												
Y	US 5,652,096 A (CIMINO) 29 July 1997, columns 7 and 10.	50-64												
Y	US 5,503,980 A (CANTOR) 02 April 1996, columns 5 and 6.	51-64												
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.														
<table border="0"> <tr> <td>* Special categories of cited documents:</td> <td>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>*A* document defining the general state of the art which is not considered to be of particular relevance</td> <td>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>*B* earlier document published on or after the international filing date</td> <td>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>*A* document member of the same patent family</td> </tr> <tr> <td>*O* document referring to an oral disclosure, use, exhibition or other means</td> <td></td> </tr> <tr> <td>*P* document published prior to the international filing date but later than the priority date claimed</td> <td></td> </tr> </table>			* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	*B* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family	*O* document referring to an oral disclosure, use, exhibition or other means		*P* document published prior to the international filing date but later than the priority date claimed	
* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention													
*A* document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone													
*B* earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art													
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family													
*O* document referring to an oral disclosure, use, exhibition or other means														
*P* document published prior to the international filing date but later than the priority date claimed														
Date of the actual completion of the international search 27 OCTOBER 1998		Date of mailing of the international search report 03 NOV 1998												
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer  CARLA MYERS Telephone No. (703) 308-0196												



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US98/16966

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y, P	US 5,683,657 A (MIAN) 04 November 1997, column 7.	51-64

**INTERNATIONAL SEARCH REPORT**

International application No.

PCT/US98/16966

**B. FIELDS SEARCHED**

Electronic data bases consulted (Name of data base and where practicable terms used):

APS; DIALOG: Medline, CA, Derwent Patents, Biosis, Embase, SciSearch

search terms: probe, array, immobilize, hydrophobic, microtitre, chip, particle, bead, flow cytometry, hybridization, mismatch, formamide, betaine, spermidine, spermine, ligase, ligate

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**